A CLUB GUESSING TOOLBOX I

TANMAY INAMDAR AND ASSAF RINOT

ABSTRACT. Club guessing principles were introduced by Shelah as a weakening of Jensen's diamond. Most spectacularly, they were used to prove Shelah's ZFC bound on $2^{\aleph_{\omega}}$. These principles have found many other applications: in cardinal arithmetic and PCF theory; in the construction of combinatorial objects on uncountable cardinals such as Jónsson algebras, strong colourings, Souslin trees, and pathological graphs; to the non-existence of universals in model theory; to the non-existence of forcing axioms at higher uncountable cardinals; and many more.

In this paper, the first part of a series, we survey various forms of clubguessing that have appeared in the literature, and then systematically study the various ways in which a club-guessing sequences can be improved, especially in the way the frequency of guessing is calibrated.

We include an expository section intended for those unfamiliar with clubguessing and which can be read independently of the rest of the article.

1. INTRODUCTION

1.1. Motivation. In this paper, we initiate a study of various aspects and forms of *club guessing*. Our definitions are quite general, and in order to motivate them we start with a brief survey of the various forms of club guessing that have appeared in the literature as well as their applications. All undefined notation can be found in Section 1.4, but we remind the reader right away that for a pair $\lambda < \kappa$ of infinite regular cardinals, $E_{\lambda}^{\kappa} := \{\delta < \kappa \mid cf(\delta) = \lambda\}$, and $E_{\neq\lambda}^{\kappa} := \{\delta < \kappa \mid cf(\delta) \neq \lambda\}$.

Shortly after Jensen constructed Souslin trees in L [Jen68], he isolated a combinatorial principle named *diamond*, which is sufficient for the construction.

Fact 1.1 ([Jen72]). In Gödel's constructible universe L, for every regular uncountable cardinal κ and every stationary $S \subseteq \kappa$, there is a sequence $\vec{A} = \langle A_{\delta} \mid \delta \in S \rangle$ which is a $\Diamond(S)$ -sequence, that is:

- (i) for every $\delta \in S$, A_{δ} is a subset of δ ;
- (ii) for every subset $A \subseteq \kappa$, the set $\{\delta \in S \mid A_{\delta} = A \cap \delta\}$ is stationary in κ .

It is easy to see that for $S \subseteq \kappa$ as above, $\Diamond(S)$ implies that $2^{<\kappa} = \kappa$, and hence diamond is not a consequence of ZFC. In contrast, the following result of Shelah [She94c], which is the most well-known club-guessing result, is a theorem of ZFC.

Fact 1.2 ([She94c]). Suppose that $\lambda < \lambda^+ < \kappa$ are infinite regular cardinals. Then there is a sequence $\vec{C} = \langle C_{\delta} | \delta \in E_{\lambda}^{\kappa} \rangle$ such that

- (i) for every $\delta \in E_{\lambda}^{\kappa}$, C_{δ} is a club in δ ;
- (ii) for every club $D \subseteq \kappa$, the set $\{\delta \in S \mid C_{\delta} \subseteq D\}$ is stationary in κ .

Date: Preprint as of July 22, 2022. For the latest version, visit http://p.assafrinot.com/46.

In particular, unlike the \Diamond principle or its descendants the \clubsuit and \uparrow principles (see [Ost76, BGKT78]), the focus is not on predicting arbitrary or even just cofinal subsets of κ , but rather only the closed and unbounded subsets of κ . This makes the task of guessing easier, since the collection of club subsets of κ generate a normal κ -complete filter.

The most famous application of Fact 1.2 is Shelah's PCF bound (see [She94a] or [AM10, Theorem 7.3]):

$$2^{\aleph_{\omega}} \leq \max\{2^{\aleph_0}, \aleph_{\omega_4}\}.$$

Apart from upper bounds on cardinal exponentiation, Fact 1.2 has many other uses in PCF theory. As an example from the basic theory, in obtaining exact upper bounds for sequences of ordinal functions (see [AM10, Lemma 2.19]), in fact showing that there are stationary sets consisting of points of large cofinality in the approachability ideal $I[\lambda]$ (see [Eis10b, §3]). Outside of PCF theory, there are applications of Fact 1.2 to the universality spectrum of models [KS92, Dza05, She21], cardinal invariants of the continuum [Zap04, Zap08], cardinal invariants at uncountable cardinals [BHZ07, GHZ21], to the study of the Boolean algebra $P(\lambda)/[\lambda]^{<\lambda}$ for λ singular of countable cofinality [KS01], to showing the incompactness of chromatic number [She13], to obtain a refinement of the downwards Löwenheim-Skolem Theorem [FT05], to study the saturation of the non-stationary ideal on $P_{\lambda}(\kappa)$ [Shi99], to obtain two-cardinal diamond principles in ZFC [Tod02, Shi08, Mat09, SheXX], to obtaining consequence of forcing axioms [She08], to obtain limitative results on forcing [RaS96, Tod18], to constructing graphs with a prescribed rate of growth of the chromatic number of its finite subgraphs [LH20]. That Fact 1.2 is a theorem of ZFC also imposes important limitations on the theory of forcing axioms at successors of uncountable cardinals (see for example [She03b, She03c, Nee14]).

While club guessing was motivated by finding a weak substitute for the diamond principle, in [She10], Shelah, using arguments that materialized through the development of the theory of club guessing, proved the next theorem on diamond, concluding a 40 year old search for such a result (see the review in [Rin10b]).

Fact 1.3 ([She10]). Let λ be an uncountable cardinal, and let $S \subseteq E_{\neq cf(\lambda)}^{\lambda^+}$ be stationary. The following are equivalent:

- (1) $2^{\lambda} = \lambda^+$;
- (2) $\Diamond(S)$.

Apart from Fact 1.2, there are other, finer, forms of club guessing which are less well-known and yet altogether have a variety of applications. For instance, Fact 1.2 says nothing about the case $\kappa = \lambda^+$. For this, we have the following result of Shelah.

Fact 1.4 ([She94c, Claim 2.4]). There is a sequence $\vec{C} = \langle C_{\delta} \mid \delta \in E_{\aleph_1}^{\aleph_2} \rangle$ such that

- (i) for every $\delta \in E_{\aleph_1}^{\aleph_2}$, C_{δ} is a club in δ of ordertype ω_1 ; (ii) for every club $D \subseteq \aleph_2$, there is a $\delta \in E_{\aleph_1}^{\aleph_2}$ such that the following set is cofinal in δ :

$$\{\beta < \delta \mid \min(C_{\delta} \setminus (\beta + 1)) \in D \cap E_{\aleph_1}^{\aleph_2}\}.$$

To compare the preceding with Fact 1.2, we see two differences in the corresponding Clause (ii). The first, here we require a single δ instead of stationarily many, however, this is easily seen to be equivalent.¹ Second, which is more important, instead of requiring C_{δ} to be a subset of D, we now merely require that the intersection of D with $(E_{\aleph_1}^{\aleph_2} \text{ and})$ the set $\operatorname{nacc}(C_{\delta})$ of all non-accumulation points of C_{δ} be cofinal in δ . This choice is not arbitrary. For club many $\delta \in E_{\aleph_1}^{\aleph_2}$, both the set $\operatorname{acc}(C_{\delta})$ of all accumulation points of C_{δ} and the set $D \cap \delta$ are clubs in δ , and hence $\operatorname{acc}(C_{\delta}) \cap D$ is trivially cofinal in δ .

Consider now another example due to Shelah concerning the case $\kappa = \lambda^+$ (see [SS10], for a short proof):

Fact 1.5 ([She03a, Claim 3.5]). There is a sequence $\vec{C} = \langle C_{\delta} \mid \delta \in E_{\aleph_1}^{\aleph_2} \rangle$ such that

- (i) for every $\delta \in E_{\aleph_1}^{\aleph_2}$, C_{δ} is a club in δ of order type ω_1 ;
- (ii) for every club $D \subseteq \aleph_2$, there is a $\delta \in S$ such that the following set is stationary in δ :

$$\{\beta < \delta \mid \min(C_{\delta} \setminus (\beta + 1)) \in D\}.$$

Comparing the two, we see that Fact 1.4 features a sequence where the guessing is measured against the ideal $J^{\text{bd}}[\delta]$ of bounded subsets of δ , whereas here we measure against the nonstationary ideal NS_{δ}. However, Fact 1.4 features a sequence which guesses clubs *relative to the set* $E_{\aleph_1}^{\aleph_2}$, and for this reason the two results are incomparable. In this paper, a join of the two results is obtained.

Note that so far we have always required that the clubs C_{δ} have the minimal possible ordertype of $cf(\delta)$. The small ordertype requirement is trivially gotten for a sequence that guesses as in Clause (ii) of Fact 1.2. In other cases, however, obtaining that the local clubs have a small ordertype requires extra care (see for example [Koj95, Theorem 9], where the weaker form of Fact 1.4 is proved where one requires each of the C_{δ} to merely have size \aleph_1). As a sample application, we mention that in [HHS09], a strong form of club-guessing at \aleph_1 with minimal ordertype is used to construct a small Dowker space.

However, attention should not be restricted only to guessing with minimal ordertypes. At the level of \aleph_1 , the ordertypes of guessing sequences play a crucial role in separating forcing axioms at \aleph_1 in [She98, Chapter XVII], and later in [AFMS13] as well. At higher cardinals, guessing sequences \vec{C} with very large ordertypes are useful for getting a pathological graph $G(\vec{C})$ with maximal chromatic number [Rin15a]. An open question concerning guessing sequences of maximal ordertype is stated in [Rin14c, Question 2]. For an extended discussion, see the introduction to [Rin15b].

At a cardinal κ that is a limit or a successor of a limit, another type of relative club-guessing has shown to be useful, where the guessing feature stipulates additional conditions on the sequence $\langle cf(\gamma) | \gamma \in nacc(C_{\delta}) \rangle$. In [She94c, Hof13], the additional condition is that this sequence is strictly increasing and converging to $|\delta|$. This is used to construct colourings satisfying strong negative square bracket partition relations [ES05, ES09]. An earlier construction (see [She94c] or [Eis10b, Theorem 5.19]) requires that the sequence $\langle cf(\gamma) | \gamma \in nacc(C_{\delta}) \rangle$ have cofinally many cardinals carrying a Jónsson algebra. This is used to construct Jónsson algebras at κ . Note that the existence of a club-guessing sequence of large ordertypes in ZFC would give rise to such sequences, in particular, solving [ES09, Question 2.4] in the affirmative.

We move on to the next example, this time a question of Shelah.

¹This equivalence is true in greater generality; see the discussion at the end of Section 2.

Question 1.6 ([She00, Question 5.4]). Let $\lambda < \lambda^+ = \kappa$ be regular uncountable cardinals. Is there a sequence $\vec{C} = \langle C_{\delta} | \delta \in E_{\lambda}^{\kappa} \rangle$ such that

- (i) for every $\delta \in E_{\lambda}^{\kappa}$, C_{δ} is a club in δ of ordertype λ ;
- (ii) for every club $D \subseteq \kappa$, there is a $\delta \in E_{\lambda}^{\kappa}$ such that the following set is stationary in δ :

 $\{\beta < \delta \mid \beta_1, \beta_2 \in D \text{ where } \beta_1 := \min(C_{\delta} \setminus (\beta + 1)) \& \beta_2 := \min(C_{\delta} \setminus (\beta_1 + 1))\}.$

Compared to Fact 1.5, here we require that for stationarily many $\beta < \delta$, two consecutive non-accumulation points following β are in the club D. Shelah mentions (without proof) that this slight strenghening of Fact 1.5 combined with GCH allows for the construction of a κ -Souslin tree. This is related to the open problem of whether GCH implies the existence of an \aleph_2 -Souslin tree (see [Rin19]), and the earlier work of Kojman and Shelah on that matter [KS93].

Here we shall prove that at the level of \aleph_2 , an affirmative answer to Shelah's question follows from the existence of a sequence as in Fact 1.4 in which $D \cap E_{\aleph_1}^{\aleph_2}$ in Clause (ii) is replaced by $D \cap E_{\aleph_0}^{\aleph_2}$. However, Asperó has answered Shelah's question negatively [Asp14]. Getting this failure together with the GCH remains open.

The feature of guessing consecutive points has other applications in the construction of Souslin trees: in [BR21, §5], the feature of guessing with two consecutive points allows to reduce a $\Diamond(\kappa)$ hypothesis from [BR17a] to just $\kappa^{<\kappa} = \kappa$. In [BR17b], a feature of guessing with ω -many consecutive points is used to construct Souslin trees with precise control over their reduced powers.

Returning to the discussion after Fact 1.5, there is another way to impose that the set of good guesses be 'large'. Here is an example, again due to Shelah.

Fact 1.7 ([She97, Claim 3.10]). Suppose that $\kappa = \lambda^+$ for a regular uncountable cardinal λ that is not strongly inaccessible. Then, there is a sequence $\langle h_{\delta} : C_{\delta} \to \omega | \delta \in E_{\lambda}^{\kappa} \rangle$ such that

- (1) for every $\delta \in E_{\lambda}^{\kappa}$, C_{δ} is a club in δ of ordertype λ ;
- (2) for every club $D \subseteq \kappa$, there is a $\delta \in E_{\lambda}^{\kappa}$ such that

$$\bigwedge_{n < \omega} \sup\{\beta < \delta \mid \min(C_{\delta} \setminus (\beta + 1)) \in D \cap h_{\delta}^{-1}\{n\}\} = \delta.$$

This result is used in [Rin14b, §2] in producing a strong oscillation with ω many colours, sufficient to derive strong colorings [Rin14b, §3] and transformations of the transfinite plane [RZ21]. Any improvement of the above result that partitions the club-guessing into θ many pieces immediately translates to getting a strong oscillation with θ colours. This also connects to our previous discussion on guessing sequences $\langle C_{\delta} | \delta \in S \rangle$ with very large ordertypes, since the number of pieces into which C_{δ} may be partitioned is bounded by $|C_{\delta}|$.

We shall later show that for the purpose of obtaining such partitioned clubguessings, the move from the unbounded ideal to the non-stationary ideal as in Fact 1.4 to Fact 1.5 is beneficial. Sufficient conditions and applications to an even stronger form of partitioned club-guessing in which there is global function h: $\kappa \to \theta$ such that $h_{\delta} = h \upharpoonright C_{\delta}$ for all δ may be found in [She94c, §3] and [LHR18, Theorem 4.20].

A very useful feature of club-guessing sequences we have so far ignored is *coher*ence. Coherent club-guessing sequences have been applied to set theory of the real line [BLS17], and to cardinal invariants of the continuum [SZ02]. Coherent clubguessing sequences were also used to show the non-existence of a natural forcing axiom [She04] and to construct strong colourings [Rin14a, RZ22]. Weakly coherent club-guessing at the level of \aleph_1 have been used to define a pathological topology on the real line [Zap01], and weakly coherent club-guessing at the level of a successor λ^+ of a singular cardinal λ was used in [BR19a, §2.1] to prevent λ -distributive λ^+ trees from having a cofinal branch, thus, yielding nonspecial λ^+ -Aronszajn trees.

The above is hardly an exhaustive list of applications of club guessing but merely a selection biased by the themes of this paper. Additional key results, including those from [GS97, Ish05, FK05, Eis10a] will be discussed in Part II of this series. Another caveat is that in this paper we shall only be concerned with getting clubguessing results at $\kappa \geq \aleph_2$. The behavior of club-guessing at the level of \aleph_1 is entirely independent of ZFC, and we refer the reader to [Hir07, Moo08, EN09, IL12a, IL12b, Ish15, AK20, GS22] for more on that matter.

1.2. The results. Throughout the paper, κ stands for an arbitrary regular uncountable cardinal; θ, μ, χ are (possibly finite) cardinals $\leq \kappa, \lambda$ and ν are infinite cardinals $< \kappa, \xi, \sigma$ are nonzero ordinals $\leq \kappa$, and S and T are stationary subsets of κ . We shall sometimes implicitly assume that S consists of nonzero limit ordinals.

Definition 1.8. A *C*-sequence over *S* is a sequence $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ such that, for every $\delta \in S$, C_{δ} is a closed subset of δ with $\sup(C_{\delta}) = \sup(\delta)$. It is said to be ξ -bounded if $\operatorname{otp}(C_{\delta}) \leq \xi$ for all $\delta \in S$.

Our first main result fulfills the promise of finding a join of Facts 1.4 and 1.5.

Theorem A. For every successor cardinal λ , there exists a λ -bounded C-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in E_{\lambda}^{\lambda^{+}} \rangle$ satisfying the following. For every club $D \subseteq \lambda^{+}$, there is a $\delta \in E_{\lambda}^{\lambda^{+}}$ such that the following set is stationary in δ :

$$\{\beta < \delta \mid \min(C_{\delta} \setminus (\beta + 1)) \in D \cap E_{\lambda}^{\lambda^{+}}\}.$$

Our next result deals with coherent guessing sequences.

Theorem B. For every cardinal $\lambda \geq \beth_{\omega}$ such that $\square(\lambda^+)$ holds, for all stationary subsets S, T of λ^+ , there exists an \sqsubseteq^* -coherent C-sequence $\vec{C} = \langle C_{\delta} | \delta < \lambda^+ \rangle$ such that, for every club $D \subseteq \lambda^+$, there is a $\delta \in S$ such that $\sup(\operatorname{nacc}(C_{\delta}) \cap D \cap T) = \delta$.

The next two results address the problem of partitioning a given club-guessing sequence into θ many pieces as in Fact 1.7.²

Theorem C. Suppose that for each $\delta \in E_{\lambda}^{\kappa}$, J_{δ} is some λ -complete ideal over δ , and suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in E_{\lambda}^{\kappa} \rangle$ is a given λ -bounded C-sequence satisfying that for every club $D \subseteq \lambda$, there exists $\delta \in E_{\lambda}^{\kappa}$ such that

$$\{\beta < \delta \mid \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T\} \in J_{\delta}^+.$$

Any of the following hypotheses imply that there exists a map $h : \lambda \to \theta$ such that for every club $D \subseteq \lambda$, there is a $\delta \in E_{\lambda}^{\kappa}$ such that, for every $\tau < \theta$,

$$\{\beta < \delta \mid h(\operatorname{otp}(C_{\delta} \cap \beta)) = \tau \& \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T\} \in J_{\delta}^{+}$$

 $^{^{2}}$ The related problem of using a given club-guessing sequence to produce another club-guessing sequence that admits a partition is also addressed in this paper. See Theorem 4.12 and the introduction to Section 6.

(i) $\theta = \lambda = \lambda^{<\lambda}$ and λ is a successor cardinal; (ii) $\theta = \lambda$, $\Diamond(\lambda)$ holds, and λ is not Mahlo;

(iii) $\theta = \lambda$, $\diamondsuit^*(\lambda)$ holds, and each J_{δ} is normal;

(iv) $\theta < \lambda$ is regular, and λ is not greatly Mahlo.

Theorem D. Under the same setup of the previous theorem, any of the following hypotheses imply that there exists a sequence of maps $\langle h_{\delta} : \lambda \to \theta \mid \delta \in E_{\lambda}^{\kappa} \rangle$ such that for every club $D \subseteq \lambda$, there is a $\delta \in E_{\lambda}^{\kappa}$ such that, for every $\tau < \theta$,

$$\{\beta < \delta \mid h_{\delta}(\operatorname{otp}(C_{\delta} \cap \beta)) = \tau \& \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T\} \in J_{\delta}^{+}.$$

(i) $\theta = \lambda$ and $\diamondsuit^*(\lambda)$ holds;

(ii) λ is not strongly inaccessible, and θ is the least to satisfy $2^{\theta} \geq \lambda$;

(*iii*) $\theta^+ = \lambda;$

(iv) $\theta = \omega$ and λ is not ineffable.

Our last result fulfills the promise to show that at the level of \aleph_2 an affirmative answer to Question 1.6 follows from the existence of a sequence as in Fact 1.4 in which $D \cap E_{\aleph_1}^{\aleph_2}$ in Clause (ii) is replaced by $D \cap E_{\aleph_0}^{\aleph_2}$.

Theorem E. For every successor cardinal λ , if there exists a λ -bounded C-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in E_{\lambda}^{\lambda^+} \rangle$ such that for every club $D \subseteq \lambda^+$, there is a $\delta \in E_{\lambda}^{\lambda^+}$ such that $\sup(\operatorname{nacc}(C_{\delta}) \cap D \cap E_{<\lambda}^{\lambda^+}) = \delta$, then there exists a λ -bounded C-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in E_{\lambda}^{\lambda^+} \rangle$ such that for every club $D \subseteq \lambda^+$, there is a $\delta \in E_{\lambda}^{\lambda^+}$ such that the following set is stationary in δ :

 $\{\beta < \delta \mid \beta_1, \beta_2 \in D \text{ where } \beta_1 := \min(C_\delta \setminus (\beta + 1)) \& \beta_2 := \min(C_\delta \setminus (\beta_1 + 1))\}.$

1.3. Organization of this paper. The aim of Section 2 is to give the reader a tour of the basic methods for proving club-guessing theorems. The purpose is introductory, and with one exception, all the results we prove are not new, though some of them are not widely known. In particular, we give a proof of Fact 1.2. In Subsection 2.1 our main definition, Definition 2.2, can be found.

In Section 3, our theme is to obtain club-guessing sequences with additional *coherence* properties. This is done by starting with an arbitrary *C*-sequence with some degree of coherence, and then improving it to make it guess clubs as well, all the while preserving the coherence. This allows us to obtain 'coherent forms' of known results such as Fact 1.4. At the end of the section, we record the results which can be obtained by the same proofs, but without any assumptions of coherence on the initial *C*-sequence. A proof of Theorem B can be found in this section.

In Section 4, we consider partitioned club-guessing. We show how the colouring principles of [IR22a, IR22b] allow us to not just obtain *partitioned* club guessing, but in fact *partition* club guessing (recall Footnote 2). Furthermore, using these colouring principles allows us to separate the combinatorial content from the club guessing content in previous results about partitioned club guessing (see [She97, Lemma 3.10]). A proof of Theorems C and D can be found here.

In Section 5, we turn to the problem of guessing many consecutive non-accumulation points as in the discussion surrounding Question 1.6. We show how a sequence guessing clubs relative to points of small cofinality can be modified for this purpose.

In the last section, Section 6, our focus is on improving the quality of the guessing calibrated against the ideal. Mainly, our focus is moving from the unbounded ideal

 $\mathbf{6}$

to the non-stationary as in the move from Fact 1.4 to Fact 1.5. Similar ideas also allow us to improve some results from Section 4. The proofs of Theorems A and E can be found here.

1.4. Notation and conventions. We have already listed some conventions in the beginning of Subsection 1.2. Here, we list some more. Let $\log_{\chi}(\lambda)$ denote the least cardinal $\theta \leq \lambda$ such that $\chi^{\theta} \geq \lambda$. For sets of ordinals A, B, we denote $A \circledast B := \{(\alpha, \beta) \in A \times B \mid \alpha < \beta\}$ and we identify $[B]^2$ with $B \circledast B$. For $\theta > 2$, $[\kappa]^{\theta}$ simply stands for the collection of all subsets of κ of size θ . For sets of ordinals A, B, we write $A \sqsubseteq B$ iff there exists an ordinal δ such that $A = B \cap \delta$; we write $A \sqsubseteq^* B$ iff there exists a pair of ordinals $\epsilon < \delta$ such that $A \setminus \epsilon = B \cap [\epsilon, \delta)$.

Let $E_{\theta}^{\kappa} := \{ \alpha < \kappa \mid \mathrm{cf}(\alpha) = \theta \}$, and define $E_{\leq \theta}^{\kappa}$, $E_{\geq \theta}^{\kappa}$, $E_{\geq \theta}^{\kappa}$, $E_{\geq \theta}^{\kappa}$, $E_{\neq \theta}^{\kappa}$ analogously. For a stationary $S \subseteq \kappa$, we write $\mathrm{Tr}(S) := \{ \alpha \in E_{\geq \omega}^{\kappa} \mid S \cap \alpha \text{ is stationary in } \alpha \}$.

For a set of ordinals A, we write $\operatorname{ssup}(A) := \operatorname{sup}\{\alpha + 1 \mid \alpha \in A\}$, $\operatorname{acc}^+(A) := \{\alpha < \operatorname{ssup}(A) \mid \operatorname{sup}(A \cap \alpha) = \alpha > 0\}$, $\operatorname{acc}(A) := A \cap \operatorname{acc}^+(A)$, $\operatorname{nacc}(A) := A \setminus \operatorname{acc}(A)$, and $\operatorname{cl}(A) := A \cup \operatorname{acc}^+(A)$. A function $f : A \to \operatorname{Ord}$ is regressive iff $f(\alpha) < \alpha$ for every nonzero $\alpha \in A$. A function $f : [A]^2 \to \operatorname{Ord}$ is upper-regressive iff $f(\alpha, \beta) < \beta$ for every pair $(\alpha, \beta) \in [A]^2$.

2. WARMING UP

In this introductory section we have two tasks. The first is to introduce our main definition, Definition 2.2, and the second is to familiarise the reader with the basic idea of all club-guessing proofs, the method of 'collecting counterexamples'.³ The former is achieved in Subsection 2.1.

For the latter purpose, we provide in Subsection 2.2 a proof of some known clubguessing results including the most famous, Fact 1.2. In the process we hope to make the reader comfortable with Definition 2.2. However, we stress that the full generality of Definition 2.2 is not needed in Subsection 2.2.

2.1. **Preliminaries.** The aim of our main definition, Definition 2.2, is to provide a language that is able to differentiate between all of the club-guessing principles we have met in Subsection 1.1. While there are numerous parameters in the definition, we hope to have convinced the reader with the examples from Section 1.1 that all of them have been found fruitful from the point of view of applications.

Definition 2.1 ([BR17a]). For a set of ordinals C, write

$$\operatorname{succ}_{\sigma}(C) := \{ C(j+1) \mid j < \sigma \& j+1 < \operatorname{otp}(C) \}.$$

In particular, for all $\gamma \in C$ such that $\sup(\operatorname{otp}(C \setminus \gamma)) \geq \sigma$, $\operatorname{succ}_{\sigma}(C \setminus \gamma)$ consists of the next σ -many successor elements of C above γ .

Throughout the paper, we shall be working with some sequence $\vec{J} = \langle J_{\delta} | \delta \in S \rangle$ such that, for each $\delta \in S$, J_{δ} is a cf(δ)-additive proper ideal over δ extending $J^{\text{bd}}[\delta] := \{B \subseteq \delta \mid \sup(B) < \delta\}.$

³As Shelah puts it in [She94c]: "The moral is quite old-fashioned: if you work hard and continue to try enough times, correcting and recorrecting yourself you will eventually succeed."

Definition 2.2 (Main definition). $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$ asserts the existence of a ξ bounded *C*-sequence $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ such that, for every club $D \subseteq \kappa$ there is a $\delta \in S$ such that

 $\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} \in J_{\delta}^+.$

Convention 2.3. We will often simplify the notation by omitting some parameters, in which case, these parameters take their weakest possible values. Specifically, if we omit ξ , then we mean that $\xi = \kappa$; if we omit \vec{J} , then we mean that $\vec{J} = \langle J^{\mathrm{bd}}[\delta] \mid \delta \in S \rangle$, if we omit \vec{J} and σ , then we mean that $\sigma = 1$ and $\vec{J} = \langle J^{\mathrm{bd}}[\delta] \mid \delta \in S \rangle$.

The following propositions collect some evident properties of CG.

Proposition 2.4 (Monotonicity). Suppose that $CG_{\xi}(S, T, \sigma, \langle J_{\delta} | \delta \in S \rangle)$ holds, as witnessed by a sequence \vec{C} . Then, assuming all of the following conditions are satisfied, \vec{C} also witnesses that $CG_{\xi'}(S', T', \sigma', \langle J'_{\delta} | \delta \in S' \rangle)$ holds.

(i) $\xi \leq \xi'$; (ii) $S \subseteq S'$; (iii) $T \subseteq T'$; (iv) $\sigma \geq \sigma'$; (v) for each $\delta \in S'$, $J'_{\delta} \supseteq J_{\delta}$.

Proposition 2.5 (Indecomposability). Suppose that $CG_{\xi}(S, T, \sigma, \vec{J})$ holds, as witnessed by a sequence \vec{C} .

- (1) For every regressive map $f: S \to \kappa$, there exists some $i < \kappa$ such that \vec{C} restricted to $S_i := f^{-1}\{i\}$ witnesses $\operatorname{CG}_{\mathcal{E}}(S_i, T, \sigma, \vec{J})$;
- (2) For every decomposition $T = \biguplus_{i < \theta} T_i$, if $S \subseteq E_{>\theta}^{\kappa}$, then there exists some $i < \theta$ such that \vec{C} witnesses $\operatorname{CG}_{\mathcal{E}}(S, T_i, \sigma, \vec{J})$.

2.2. A tour of club-guessing. Most of the results in this article will have the following format: we shall assume the existence of a C-sequence \vec{C} witnessing a certain form of club guessing, and then we shall improve or modify this \vec{C} so that it satisfies another form of club guessing, or such that it has some other properties. For example, Proposition 2.4 suggests to us that starting from \vec{C} witnessing $\operatorname{CG}_{\xi}(S,T,\sigma,\langle J_{\delta} \mid \delta \in S \rangle)$, we may look into the possibility of reducing ξ , or shrinking S or T, or increasing σ , or enlarging the ideals in $\langle J_{\delta} \mid \delta \in S \rangle$, all the while preserving the guessing properties of \vec{C} . We shall be considering these problems and other similar ones in this article.

What is important in all this is our ability to modify a given C-sequence to satisfy other, or additional, properties. In this section we shall present some of the standard techniques that one uses to make such modifications, and we do this by giving a proof of Fact 1.2 (Corollary 2.14 below). As our purpose is introductory, we avoid giving the most direct proofs and focus instead upon the gradual process of improving the guessing.

We then move on to proving a less-known theorem of Shelah that $CG(S, \kappa)$ holds for every stationary subset S of every regular cardinal $\kappa \geq \aleph_2$ (Theorem 2.15 below).

We finish by giving in Proposition 2.22 an example of how a prediction principle weaker than \diamondsuit consisting of a matrix of sets can be modified to obtain a club-guessing principle.

We begin by considering a very weak variation of $CG_{\mathcal{E}}(S,T)$.

Definition 2.6. $\operatorname{CG}_{\xi}(S, T, -)$ asserts the existence of a ξ -bounded *C*-sequence, $\langle C_{\delta} \mid \delta \in S \rangle$ such that, for every club $D \subseteq \kappa$ there is a $\delta \in S$ with $\sup(C_{\delta} \cap D \cap T) = \delta$.

The following might be obvious, but since we have just begun, we give a detailed proof.

Proposition 2.7. For every triple of regular cardinals $\mu < \lambda < \kappa$, for every stationary $S \subseteq E_{\lambda}^{\kappa}$, any λ -bounded C-sequence over S witnesses $CG_{\lambda}(S, E_{\mu}^{\kappa}, -)$.

Proof. Let $S \subseteq E_{\lambda}^{\kappa}$ be stationary, and let $\langle C_{\delta} \mid \delta \in S \rangle$ be a λ -bounded C-sequence. Given a club D in κ , pick $\delta \in S \cap \operatorname{acc}(D)$. Since $\operatorname{cf}(\delta) = \lambda$ which is a regular uncountable cardinal, $D \cap \delta$ is club in δ . Pick a closed and unbounded subset B_{δ} of δ of ordertype λ such that $B_{\delta} \subseteq D$. Since C_{δ} is also club in δ , and δ has uncountable cofinality, $B_{\delta} \cap C_{\delta}$ is as well club in δ . Let $\langle a_{\delta}(i) \mid i < \lambda \rangle$ be the increasing enumeration of $C_{\delta} \cap D_{\delta}$. As this is an increasing and continuous sequence, it is clear then that for every $j \in \operatorname{acc}(\lambda)$, $\operatorname{cf}(a_{\delta}(j)) = \operatorname{cf}(j)$. Since $\mu < \lambda$, the set E_{μ}^{λ} is cofinal in λ , and so for every $j \in E_{\mu}^{\lambda}$, $a_{\delta}(j) \in E_{\mu}^{\kappa}$. It follows that $\{a_{\delta}(j) \mid j \in E_{\mu}^{\lambda}\}$ is a subset of $C_{\delta} \cap D \cap E_{\mu}^{\lambda}$ which is unbounded in δ .

Our goal now is to show that if $\lambda^+ < \kappa$, then $\operatorname{CG}_{\lambda}(S,T,-)$ implies that there is a *C*-sequence $\langle C_{\delta} \mid \delta \in S \rangle$ with the property that for every club $D \subseteq \kappa$, the set $\{\delta \in S \mid C_{\delta} \subseteq D\}$ is stationary. In doing so, the challenge lies in improving instances of " $\sup(C_{\delta} \cap D) = \delta$ " into instances of " $C_{\delta} \subseteq D$ ". A natural approach is to shrink each club C_{δ} into a smaller club in δ , say $\Phi(C_{\delta})$. In an ideal scenario, a single such act of shrinking will be enough and we will have our result. If the scenario is not so ideal, we would hope that $\Phi(C_{\delta})$ is at least 'better' than C_{δ} , or 'takes care of the requirements imposed by more clubs' than C_{δ} (we will be more precise momentarily). A common strategy in club guessing is to assume that there are no such ideal scenarios, and then in this case perform this shrinking process (equivalently, improvement process) iteratively for long enough that a contradiction results.

We return to precision. We shall need the following operator in what follows for purposes we have already hinted at.

Definition 2.8. For a subset $B \subseteq \kappa$, we define the operator $\Phi^B : \mathcal{P}(\kappa) \to \mathcal{P}(\kappa)$ by letting for all $x \subseteq \kappa$,

$$\Phi^B(x) := \begin{cases} \operatorname{cl}(x \cap B), & \text{if } \sup(x \cap B) = \sup(x); \\ x \setminus \sup(x \cap B), & \text{otherwise.} \end{cases}$$

We list a few useful properties of Φ^B :

- (i) $\sup(\Phi^B(x)) = \sup(x);$
- (ii) If $\sup(x \cap B) = \sup(x)$, then $\operatorname{nacc}(\Phi^B(x))$ is a cofinal subset of $x \cap B$;

(iii) If x is a closed subset of $\sup(x)$, then $\Phi^B(x) \subseteq x$ and $\operatorname{otp}(\Phi^B(x)) \leq \operatorname{otp}(x)$.

Lemma 2.9. Suppose that $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ witnesses $\operatorname{CG}_{\lambda}(S, T, -)$, where S and T are stationary subsets of κ . If $\lambda^+ < \kappa$, then there exists a club $D \subseteq \kappa$ such that $\langle \Phi^{D \cap T}(C_{\delta}) | \delta \in S \rangle$ witnesses $\operatorname{CG}_{\lambda}(S, T, \kappa)$.

Proof. Without loss of generality, $S \subseteq \operatorname{acc}(\kappa)$. Suppose that the conclusion does not hold. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that, for

every $\delta \in S$,

$$\sup(\operatorname{nacc}(\Phi^{D\cap T}(C_{\delta})) \setminus (F^{D} \cap T)) = \delta.$$

Here, we use that

$$\operatorname{otp}(\operatorname{nacc}(\Phi^{D\cap T}(C_{\delta}))) \leq \operatorname{otp}(\Phi^{D\cap T}(C_{\delta})) \leq \operatorname{otp}(C_{\delta}) \leq \lambda < \kappa.$$

We construct now a \subseteq -decreasing sequence $\langle D_i \mid i \leq \lambda^+ \rangle$ of clubs in κ as follows:

- (i) $D_0 := \kappa;$
- (ii) $D_{i+1} := D_i \cap F^{D_i};$

(iii) for $i \in \operatorname{acc}(\lambda^+ + 1)$, $D_i := \bigcap_{i' < i} D_{i'}$.

Since $\lambda^+ < \kappa$, all these are club subsets of κ . As \vec{C} witnesses $\operatorname{CG}_{\lambda}(S, T, -)$, let us now pick $\delta \in S$ with $\sup(C_{\delta} \cap D_{\lambda^+} \cap T) = \delta$. In particular, for all $i < \lambda^+$, $\sup(C_{\delta} \cap D_i \cap T) = \sup(C_{\delta})$, so that $\Phi^{D_i \cap T}(C_{\delta}) = \operatorname{cl}(C_{\delta} \cap D_i \cap T)$. Now, as $\langle D_i |$ $i < \lambda^+ \rangle$ is \subseteq -decreasing, so is $\langle C_{\delta} \cap D_i \cap T | i < \lambda^+ \rangle$. But $\operatorname{otp}(C_{\delta}) \leq \lambda$, so we may find some $i < \lambda^+$ such that $C_{\delta} \cap D_i \cap T = C_{\delta} \cap D_{i+1} \cap T$.

By the choice of F^{D_i} , and since $D_{i+1} \subseteq F^{D_i}$, we have that

$$\sup(\operatorname{nacc}(\Phi^{D_i\cap T}(C_{\delta}))\setminus (D_{i+1}\cap T))=\delta.$$

However, $\operatorname{nacc}(\Phi^{D_i \cap T}(C_{\delta})) \subseteq C_{\delta} \cap D_i \cap T = C_{\delta} \cap D_{i+1} \cap T$, which is a contradiction.

So $\operatorname{CG}_{\lambda}(S,T,-)$ implies $\operatorname{CG}_{\lambda}(S,T,\kappa)$, provided that $\lambda^+ < \kappa$. Likewise, $\operatorname{CG}(S,T,\kappa)$ holds whenever there is a witness $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ to $\operatorname{CG}(S,T)$ such that $|C_{\delta}| < \delta$ for club many $\delta \in S$.

The instance $CG(S, T, \sigma)$ with $\sigma = \kappa$ is sometimes dubbed *tail club guessing*. The next lemma derives a stronger form of guessing from tail club guessing.

Lemma 2.10. $CG(S,T,\kappa)$ holds iff there is a C-sequence $\langle C_{\delta} | \delta \in S \rangle$ such that:

- (i) for every $\delta \in S$, $\operatorname{otp}(C_{\delta}) = \operatorname{cf}(\delta)$;
- (ii) for every club $D \subseteq \kappa$, the set $\{\delta \in S \mid \operatorname{nacc}(C_{\delta}) \subseteq D \cap T\}$ is stationary.

Proof. Only the forward implication requires an argument. Let $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ be a CG (S, T, κ) -sequence. For every $i < \kappa$, we define the operator $\Phi_i : \mathcal{P}(\kappa) \to \mathcal{P}(\kappa)$ by letting for all $x \subseteq \kappa$,

$$\Phi_i(x) := \begin{cases} x, & \text{if } x \subseteq i; \\ x \setminus i, & \text{otherwise} \end{cases}$$

It is clear that $\Phi_i(x)$ is a cofinal subset of x, and $\operatorname{nacc}(\Phi_i(x)) \subseteq \operatorname{nacc}(x)$. Furthermore, if x is club in its supremum, then so is $\Phi_i(x)$.

Claim 2.10.1. There exists $i < \kappa$ such that, for every club $D \subseteq \kappa$, the set $\{\delta \in S \mid nacc(\Phi_i(C_{\delta})) \subseteq D \cap T\}$ is stationary.

Proof. Suppose not. For each $i < \kappa$, fix a sparse enough club $D_i \subseteq \kappa$ for which $\{\delta \in S \mid \operatorname{nacc}(\Phi_i(C_{\delta})) \subseteq D_i \cap T\}$ is disjoint from D_i . Let $D := \bigwedge_{i < \kappa} D_i$. By the choice of \vec{C} , there are $\delta \in S \cap D$ and $\beta < \delta$ such that $\operatorname{succ}_{\kappa}(C_{\delta} \setminus \beta) \subseteq D \cap T$. As $\operatorname{otp}(C_{\delta}) < \kappa$, we can find an $i < \delta$ such that $\operatorname{nacc}(C_{\delta} \setminus i \subseteq D \cap T$. Then $\operatorname{nacc}(\Phi_i(C_{\delta})) \subseteq D \cap T$. But $i < \delta$ and $\delta \in D$, so that $\delta \in D_i$. This is a contradiction.

Let *i* be given by the preceding claim. The sequence $\langle \Phi_i(C_{\delta}) | \delta \in S \rangle$ satisfies Clause (ii) of the Lemma. In order to incorporate Clause (i), for each $\delta \in S$, we simply pick a club C_{δ}^{\bullet} in δ of ordertype cf(δ) such that $\operatorname{nacc}(C_{\delta}^{\bullet}) \subseteq \operatorname{nacc}(\Phi_i(C_{\delta}))$. Evidently, $\langle C_{\delta}^{\bullet} | \delta \in S \rangle$ is as sought. \Box

Putting everything together, we arrive at the following striking conclusion.

Corollary 2.11 ([She94c]). For every regular uncountable cardinal λ such that $\lambda^+ < \kappa$, for every stationary $S \subseteq E_{\lambda}^{\kappa}$, there is a C-sequence $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ such that the following two hold:

- (i) for every $\delta \in S$, $\operatorname{otp}(C_{\delta}) = \lambda$;
- (ii) for every club $D \subseteq \kappa$, the set $\{\delta \in S \mid C_{\delta} \subseteq D\}$ is stationary.

Proof. By Proposition 2.7, in particular, $CG_{\lambda}(S, E_{\aleph_0}^{\kappa}, -)$ holds. Then, Lemma 2.9 implies that so does $CG_{\lambda}(S, E_{\aleph_0}^{\kappa}, \kappa)$. Now, appeal to Lemma 2.10.

At this point, it is natural to ask whether it is possible to waive the uncountability hypothesis on λ in the preceding theorem. We shall show that this is indeed the case, by invoking an operation different than that of Φ^B .

Definition 2.12. For a subset $D \subseteq \kappa$, we define the operator $\Phi_D : \mathcal{P}(\kappa) \to \mathcal{P}(\kappa)$ by letting for all $x \subseteq \kappa$,

$$\Phi_D(x) := \begin{cases} \{\sup(D \cap \eta) \mid \eta \in x, \eta > \min(D)\}, & \text{if } \sup(D \cap \sup(x)) = \sup(x); \\ x \setminus \sup(D \cap \sup(x)), & \text{otherwise.} \end{cases}$$

We list a few useful properties of Φ_D :

- (i) $\sup(\Phi_D(x)) = \sup(x);$
- (ii) $\operatorname{otp}(\Phi_D(x)) \leq \operatorname{otp}(x);$
- (iii) If $\sup(D \cap \sup(x)) = \sup(x)$, then $\operatorname{acc}^+(\Phi_D(x)) \subseteq \operatorname{acc}^+(D) \cap \operatorname{acc}^+(x)$. If in addition, D is closed below $\sup(x)$, then $\Phi_D(x) \subseteq D$.

Lemma 2.13. Suppose that $\kappa \geq \aleph_2$, and that $\langle C_{\delta} \mid \delta \in S \rangle$ is an ω -bounded C-sequence over a stationary $S \subseteq E_{\aleph_0}^{\kappa}$.

Then there is a club $D \subseteq \kappa$ such that $\langle \Phi_D(C_\delta) | \delta \in S \rangle$ witnesses $CG_{\omega}(S, \kappa, \kappa)$.

Proof. Suppose not. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that, for every $\delta \in S$,

$$\sup(\Phi_D(C_\delta) \setminus F^D) = \delta.$$

Here we have used that since C_{δ} has ordertype ω , $\Phi_D(C_{\delta})$ has ordertype ω as well, and hence all of its points are nonaccumulation points.

As $\kappa > \aleph_1$, we may construct a \subseteq -decreasing sequence $\langle D_i \mid i \leq \omega_1 \rangle$ of clubs in κ as follows:

(i) $D_0 := \kappa;$

(ii)
$$D_{i+1} := D_i \cap F^{D_i}$$

(iii) for $i \in \operatorname{acc}(\omega_1 + 1)$, $D_i := \bigcap_{i' \le i} D_{i'}$.

Pick $\delta \in S \cap \operatorname{acc}(D_{\omega_1})$. For each $i < \omega_1$, since $D_i \cap \delta$ is a closed unbounded subset of δ , it is the case that $\Phi_{D_i}(C_{\delta}) = \{ \sup(D_i \cap \eta) \mid \eta \in C_{\delta}, \eta > \min(D_i) \}$, and $\Phi_{D_i}(C_{\delta}) \subseteq D_i$.

As $\langle D_i | i \leq \omega_1 \rangle$ is \subseteq -decreasing, for each $\eta \in C_{\delta}$, $\langle \sup(D_i \cap \eta) | i < \omega_1 \rangle$ is a weakly decreasing sequence of ordinals. By well-foundedness of the ordinals, for each $\eta \in C_{\delta}$, there must be some $i_{\eta} < \omega_1$ such that $\sup(D_i \cap \eta) = \sup(D_j \cap \eta)$ whenever $i_{\eta} \leq i < j < \omega_1$. Let $i^* := \sup_{\eta \in C_{\delta}} i_{\eta}$, which is a countable ordinal as C_{δ} is a countable set. It follows that for any $i \in [i^*, \omega_1)$, $\Phi_{D_i}(C_{\delta}) = \Phi_{D_{i+1}}(C_{\delta})$. However, $\Phi_{D_{i+1}}(C_{\delta}) \subseteq D_{i+1} \subseteq F^{D_i}$, contradicting the choice of F^{D_i} .

Putting everything together:

Corollary 2.14 ([She94c]). For every pair of infinite regular cardinals $\lambda < \kappa$ and every stationary $S \subseteq E_{\lambda}^{\kappa}$, if $\lambda^+ < \kappa$, then there is a *C*-sequence $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ with the property that for every club $D \subseteq \kappa$, the set $\{\delta \in S \mid C_{\delta} \subseteq D\}$ is stationary.

We now move on to prove a lesser-known theorem of Shelah concerning clubguessing. Unlike the previous result, in the following, S is not assumed to be a subset of E_{λ}^{κ} for some fixed cardinal $\lambda < \kappa$. So, for instance, S could be the set of regular cardinals below a Mahlo cardinal κ .

Theorem 2.15 (Shelah). Suppose $\kappa \geq \aleph_2$. For every stationary $S \subseteq \kappa$, CG $(S, \kappa, 1)$ holds.

Our proof of Theorem 2.15 goes through the notion of an amenable *C*-sequence, which is a strengthening of $\otimes_{\vec{C}}$ from [She94c, p. 134].

Definition 2.16 ([BR19a, Definition 1.3]). For a stationary $S \subseteq \kappa$, a *C*-sequence $\langle C_{\delta} | \delta \in S \rangle$ is *amenable* iff for every club $D \subseteq \kappa$, the set $\{\delta \in S | \sup(D \cap \delta \setminus C_{\delta}) < \delta\}$ is nonstationary in κ .

Fact 2.17 ([IR22a, Corollary 3.11]). For every stationary $S \subseteq \kappa$, there exists a stationary $S' \subseteq S$ such that S' carries an amenable C-sequence.

Lemma 2.18. Suppose that $S \subseteq \kappa$ is stationary and $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ is an amenable *C*-sequence. If $\kappa \geq \aleph_2$, then there exists a club $D \subseteq \kappa$ for which $\langle \Phi_D(C_{\delta}) \mid \delta \in S \rangle$ witnesses $CG(S, \kappa, 1)$.

Proof. Suppose not. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that, for every $\delta \in S$,

$$\sup(\operatorname{nacc}(\Phi_D(C_{\delta})) \cap F^D) < \delta$$

As $\kappa \geq \aleph_2$, we may construct a \subseteq -decreasing sequence $\langle D_i \mid i \leq \omega_1 \rangle$ of clubs in κ as follows:

- (i) $D_0 := \kappa;$
- (ii) $D_{i+1} := D_i \cap F^{D_i};$

(iii) for $i \in \operatorname{acc}(\omega_1 + 1)$, $D_i := \bigcap_{i' < i} D_{i'}$.

As \tilde{C} is amenable and D_{ω_1} is club in κ , we may pick some $\delta \in S$ such that $\sup(D_{\omega_1} \cap \delta \setminus C_{\delta}) = \delta$. For each $i < \omega_1$, since $D_i \cap \delta$ is a closed unbounded subset of δ , it is the case that

$$\Phi_{D_i}(C_{\delta}) = \{ \sup(D_i \cap \eta) \mid \eta \in C_{\delta}, \eta > \min(D_i) \}.$$

So $\Phi_{D_i}(C_{\delta}) \subseteq D_i$ and $\operatorname{acc}(\Phi_{D_i}(C_{\delta})) \subseteq \operatorname{acc}(D_i) \cap \operatorname{acc}(C_{\delta})$.

In addition, for each $i < \omega_1$, since $D_{i+1} \subseteq F^{D_i}$, the following ordinal is smaller than δ :

$$\epsilon_i := \sup(\operatorname{nacc}(\Phi_{D_i}(C_{\delta})) \cap D_{i+1}).$$

Claim 2.18.1. There exists $I \subseteq \omega_1$ of order type ω such that $\sup\{\epsilon_i \mid i \in I\} < \delta$.

Proof. If $cf(\delta) > \omega_1$, then just let $I := \omega$. If $cf(\delta) = \omega$, then pick a countable cofinal subset E of δ and for each $i \in \omega_1$, find the least $\epsilon \in E$ such $\epsilon_i \leq \epsilon$. By the pigeonhole principle, there is an $\epsilon \in E$ for which $\{i \in I \mid \epsilon_i \leq \epsilon\}$ is uncountable. In particular, this set contains a subset of ordertype ω .

Fix I as in the claim, and then pick $\gamma \in D_{\omega_1} \cap \delta \setminus C_{\delta}$ above $\sup \{ \epsilon_i \mid i \in I \}$. As $\gamma \notin C_{\delta}$, $\eta := \min(C_{\delta} \setminus \gamma)$ is in $\operatorname{nacc}(C_{\delta})$. As $\langle \sup(D_i \cap \eta) \mid i \in I \rangle$ is a weakly decreasing sequence of ordinals, by well-foundedness there must be a pair of ordinals i < j in I such that $\beta_i := \sup(D_i \cap \eta)$ is equal to $\beta_j := \sup(D_j \cap \eta)$.

As $\gamma \in D_{\omega_1} \subseteq D_i$, $\epsilon_i < \gamma \leq \beta_i \leq \eta$, so $\beta_i \in \Phi_{D_i}(C_{\delta}) \cap (\epsilon_i, \eta]$. Likewise, $\beta_j \in \Phi_{D_i}(C_{\delta}) \cap (\epsilon_j, \eta]$. Recalling that $\beta_i = \beta_j \in D_j \subseteq D_{i+1}$, it follows that β_i is an element of $\Phi_{D_i}(C_{\delta}) \cap D_{i+1}$ above ϵ_i and hence $\beta_i \in \operatorname{acc}(\Phi_{D_i}(C_{\delta}))$. However, $\operatorname{acc}(\Phi_{D_i}(C_{\delta})) \subseteq \operatorname{acc}(D_i) \cap \operatorname{acc}(C_{\delta})$, and hence $\beta_i \in \operatorname{acc}(C_{\delta})$. But $\gamma \leq \beta_i \leq \eta$ and $C_{\delta} \cap [\gamma, \eta] = \{\eta\}$, and hence $\beta_i = \eta$, contradicting the fact that $\eta \in \operatorname{nacc}(C_{\delta})$.

Proof of Theorem 2.15. Given a stationary $S \subseteq \kappa$, appeal to Fact 2.17 to find an amenable C-sequence $\langle C_{\delta} \mid \delta \in S' \rangle$ for some stationary $S' \subseteq S$. Then, by Lemma 2.18, $CG(S', \kappa, 1)$ holds. So $CG(S, \kappa, 1)$ holds as well.

As pointed out in the introduction, if $\Diamond(S)$ holds for a given stationary subset S of κ , then, for every stationary $T \subseteq \kappa$, $CG(S, T, \kappa)$ holds. The next result shows how to get $CG(S,T,\kappa)$ from a principle weaker than $\Diamond(S)$ and even weaker than $\clubsuit(S)$ and of which many instances hold true in ZFC.

Definition 2.19 ([Rin10a]). For a stationary subset S of a regular uncountable cardinal $\kappa, \clubsuit^{-}(S)$ asserts the existence of a sequence $\langle \mathcal{A}_{\delta} | \delta \in S \rangle$ such that:

(i) for all δ ∈ S, A_δ ⊆ [δ]^{<|δ|} and |A_δ| ≤ |δ|;
(ii) for every cofinal Z ⊆ κ, there are δ ∈ S and A ∈ A_δ with sup(A ∩ Z) = δ.

Remark 2.20. Note that if $\clubsuit^{-}(S)$ holds, then $\{\delta \in S \mid cf(\delta) < |\delta|\}$ must be stationary.

Fact 2.21 ([Rin10a]). For an infinite cardinal λ and a stationary $S \subseteq \lambda^+$:

- If $S \cap E_{\neq cf(\lambda)}^{\lambda^+}$ is stationary, then $\clubsuit^-(S)$ holds;
- If \Box_{λ}^* holds and S reflects stationarily often, then $\clubsuit^-(S)$ holds.

In reading the statement of the next two propositions, keep in mind Lemma 2.10.

Proposition 2.22. Suppose that $\clubsuit^{-}(S)$ holds for a given stationary $S \subseteq \kappa$. Then, for every stationary $T \subseteq \kappa$, $CG(S, T, \kappa)$ holds.

Proof. Let $\langle \mathcal{A}_{\delta} \mid \delta \in S \rangle$ be a $\clubsuit^{-}(S)$ -sequence. For each $\delta \in S$, fix an enumeration $\{A_{\delta,i} \mid i < \delta\}$ of \mathcal{A}_{δ} .

Claim 2.22.1. There exists $i < \kappa$ such that, for every club $E \subseteq \kappa$, there is a $\delta \in S$ with $\sup(A_{\delta,i} \cap E \cap T) = \delta$.

Proof. Otherwise, for each $i < \kappa$, we may pick a counterexample E_i . Let $Z := T \cap$ $\Delta_{i < \kappa} E_i$. Pick $\delta \in S$ and $i < \delta$ such that $\sup(A_{\delta,i} \cap Z) = \delta$. Since $Z \cap (i, \delta) \subseteq T \cap E_i$, we have that $\sup(A_{\delta,i} \cap E_i \cap T) = \delta$. This contradicts the choice of E_i .

Fix i as given by the preceding claim, and denote $A_{\delta} := A_{\delta,i}$. We shall now make use of the operator Φ^B from Definition 2.8.

Claim 2.22.2. There exists a club $D \subseteq \kappa$ such that, for every club $E \subseteq \kappa$, there exists $\delta \in S$ with $\sup(A_{\delta}) = \delta$ and $\operatorname{nacc}(\Phi^{D \cap T}(A_{\delta})) \subset E$.

Proof. Suppose not. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that for every $\delta \in S$, either $\sup(A_{\delta}) < \delta$ or $\operatorname{nacc}(\Phi^{D \cap T}(A_{\delta})) \not\subseteq F^{D}$. We construct a \subseteq -decreasing sequence $\langle D_i \mid i < \kappa \rangle$ of clubs in κ as follows:

- $\begin{array}{ll} (\mathrm{i}) & D_0 := \kappa; \\ (\mathrm{ii}) & D_{i+1} := D_i \cap F^{D_i}; \\ (\mathrm{iii}) & \mathrm{for} \ i \in \mathrm{acc}(\kappa), \ D_i := \bigcap_{i' < i} D_{i'}. \end{array}$

Let $E := \Delta_{i < \kappa} D_i$. Pick $\delta \in S$ with $\sup(A_{\delta} \cap E \cap T) = \delta$. For every $i < \delta$, $\delta \in \operatorname{acc}^+(D_i \cap T)$ and $\operatorname{nacc}(\Phi^{D_i \cap T}(A_{\delta})) \not\subseteq D_{i+1}$, so that we may pick $\beta_i \in (A_{\delta} \cap C)$ $D_i \cap T$ $\setminus D_{i+1}$. As $|A_{\delta}| < \delta$, let us fix $i < j < \delta$ such that $\beta_i = \beta_j$. So $\beta_i \notin D_{i+1}$ while $\beta_j \in D_j \subseteq D_{i+1}$. This is a contradiction.

Let D be given by the preceding claim. For $\delta \in S$, let $C_{\delta} := \Phi^{D \cap T}(A_{\delta})$. Then $\langle C_{\delta} \mid \delta \in S \rangle$ witness $\operatorname{CG}(S, T, \kappa)$.

Corollary 2.23. For every uncountable cardinal λ , every stationary $S \subseteq E_{\neq cf(\lambda)}^{\lambda^+}$, and every stationary $T \subseteq \lambda^+$, $CG(S, T, \kappa)$ holds.

We end this section by saving a few words about the following natural generalisation of Definition 2.2.

Definition 2.24. For an ideal J_{κ} over κ , $\mathrm{CG}_{\xi}(J_{\kappa}, T, \sigma, \vec{J})$ asserts the existence of a C-sequence $\vec{C} = \langle C_{\delta} \mid \delta < \kappa \rangle$ such that $\{\delta < \kappa \mid \operatorname{otp}(C_{\delta}) > \xi\} \in J_{\kappa}$, and such that for every club $D \subseteq \kappa$,

$$\{\delta < \kappa \mid \{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} \in J_{\delta}^{+}\} \in J_{\kappa}^{+}.$$

First, the proof of Lemma 2.10 makes it clear that obtaining a single witness $\delta \in S$ to an instance of guessing a club is equivalent to obtaining stationarily-many such witnesses. More precisely, the usual principle $CG_{\xi}(S, T, \sigma, J)$ coincides with $\operatorname{CG}_{\xi}(J_{\kappa}, T, \sigma, \vec{J})$ for $J_{\kappa} := \operatorname{NS}_{\kappa} \upharpoonright S$. Furthermore, if \vec{C} witnesses $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$, then the collection of all $S' \subseteq S$ for which $\vec{C} \upharpoonright S'$ fails to witness $\operatorname{CG}_{\mathcal{E}}(S', T, \sigma, \vec{J})$ forms a κ -complete normal (and proper) ideal extending NS_{κ} | S.

Second, in any of the upcoming results that involve pumping an instance $CG_{\mathcal{E}}(S,\ldots)$ into a better instance $CG_{\bar{\mathcal{E}}}(S,\ldots)$, no new ideas would be needed in order to get the analogous result where S is replaced by an abstract κ -complete ideal J_{κ} over κ . In fact, for many of the results, letting J_{κ} be an \aleph_1 -indecomposable ideal (see for instance [Eis10a, §2]) over κ would be sufficient. For this reason we eschew the added generality of Definition 2.24 and focus on Definition 2.2.

3. Coherent sequences

Let us point out some commonalities in the proofs of Lemmas 2.9, 2.10, 2.13 and 2.18. In all of these, we started with a C-sequence, and then we improved it using some operation $\Phi: \mathcal{P}(\kappa) \to \mathcal{P}(\kappa)$. These similarities leads one to describe abstractly the class of such operations to which the examples we've met belong, in the hope that known members or properties of this class might be of assistance in future endeavours. This class has in fact already been delineated in work of Brodsky and Rinot in [BR19a], where they occurred in the work on constructing trees with prescribed properties by studying how the properties of a C-sequence affect the properties of the trees derived from walks on ordinals.

Definition 3.1 ([BR19a]). Let $\mathcal{K}(\kappa) := \{x \in \mathcal{P}(\kappa) \mid x \neq \emptyset \& \operatorname{acc}^+(x) \subseteq x \& \sup(x) \notin x\}$ be the set of all closed subsets of some nonzero limit ordinal $\leq \kappa$.

- An operator $\Phi : \mathcal{K}(\kappa) \to \mathcal{K}(\kappa)$ is a *postprocessing function* if for every $x \in \mathcal{K}(\kappa)$:
 - (i) $\Phi(x)$ is a club in $\sup(x)$;
 - (ii) $\operatorname{acc}(\Phi(x)) \subseteq \operatorname{acc}(x);$
- (iii) $\Phi(x) \cap \bar{\alpha} = \Phi(x \cap \bar{\alpha})$ for every $\bar{\alpha} \in \operatorname{acc}(\Phi(x))$.

Remark 3.2. By the first clause, $otp(\Phi(x)) \ge cf(sup(x))$, and by the second clause, $otp(\Phi(x)) \le otp(x)$.

It is easy to verify that the three operations we met in Section 2 — when their domains are restricted to $\mathcal{K}(\kappa)$ — are postprocessing functions. What's nice about postprocessing functions is that requirement (iii) implies that they maintain *coherence* features of *C*-sequences. Indeed, the theme of this section is to obtain club-guessing sequences which have additional coherence features. The particular coherence features we consider can be found in the following definition and in Definition 3.15 below.

Definition 3.3. Let $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ be a *C*-sequence.

- (i) For an infinite cardinal $\chi \leq \kappa$, we say that \vec{C} is $\chi \sqsubseteq$ -coherent iff for all $\delta \in S$ and $\bar{\delta} \in \operatorname{acc}(C_{\delta}) \cap E_{\geq \chi}^{\kappa}$, it is the case that $\bar{\delta} \in S$ and $C_{\bar{\delta}} = C_{\delta} \cap \bar{\delta}$;
- (ii) We say that \vec{C} is *coherent* iff it is $\omega \sqsubseteq$ -coherent;
- (iii) We say that \vec{C} is weakly coherent iff for every $\alpha < \kappa$, $|\{C_{\delta} \cap \alpha \mid \delta \in S\}| < \kappa$.

It is routine to verify that if $\Phi : \mathcal{K}(\kappa) \to \mathcal{K}(\kappa)$ is a postprocessing function, and $\langle C_{\delta} | \delta \in S \rangle$ satisfies any of the above coherence properties, then $\langle \Phi(C_{\delta}) | \delta \in S \rangle$ satisfies the same coherence property as well, Clause (iii) being key in the verification.

In particular, as a consequence of the use of postprocessing functions, in each of Lemmas 2.9, 2.10, 2.13, and 2.18, if we start with a C-sequence with one of the coherence properties above, the exact same proof ensures that the guessing C-sequence obtained satisfies the same coherence property.

As a concrete example, by [BR19a, Lemma 1.23], every transversal to a $\Box_{\xi}(\kappa, <\mu)$ sequence with $\xi < \kappa$ or $\mu < \kappa$ gives an amenable *C*-sequence $\langle C_{\delta} | \delta \in \operatorname{acc}(\kappa) \rangle$ which
can then be supplied to the machinery in Lemma 2.18. Indeed, in [BR19a], solving Question 16 from [Rin11] in the affirmative, a wide club guessing theorem was
proven using Lemma 2.18.⁴

Fact 3.4 ([BR19a, Lemma 2.5]). If $\Box_{\xi}(\kappa, <\mu)$ holds for a regular cardinal $\kappa \geq \aleph_2$ and a cardinal $\mu < \kappa$, then for every stationary $S \subseteq \kappa$, $\Box_{\xi}(\kappa, <\mu)$ may be witnessed by a sequence $\langle \mathcal{C}_{\delta} | \delta < \kappa \rangle$ with the added feature that for every club $D \subseteq \kappa$, there exists $\delta \in S$ such that, for every $C \in \mathcal{C}_{\delta}$, $\sup(\operatorname{nacc}(C) \cap D) = \delta$.

Recall 3.5. $\Box_{\xi}(\kappa, <\mu)$ asserts the existence of a sequence $\langle \mathcal{C}_{\alpha} \mid \alpha < \kappa \rangle$ satisfying all of the following:

⁴The statement of [BR19a, Lemma 2.5] does not mention the parameter ξ , however, its proof is an application of a postprocessing function and hence does not increase ordertypes.

- for every limit ordinal $\alpha < \kappa$, $1 < |\mathcal{C}_{\alpha}| < \mu$, and each $C \in \mathcal{C}_{\alpha}$ is club in α with $\operatorname{otp}(C) \leq \xi$;
- for every $\alpha < \kappa$, $C \in \mathcal{C}_{\alpha}$, and $\bar{\alpha} \in \operatorname{acc}(C)$, $C \cap \bar{\alpha} \in \mathcal{C}_{\bar{\alpha}}$;
- for every club D in κ , there exists some $\alpha \in \operatorname{acc}(D)$ such that $D \cap \alpha \notin \mathcal{C}_{\alpha}$.

Remark 3.6. The instance $\Box_{\kappa}(\kappa, <2)$ is better known as $\Box(\kappa)$, the instance $\Box_{\lambda}(\lambda^+, <2)$ is better known as \Box_{λ} , and the instance $\Box_{\lambda}(\lambda^+, <\lambda^+)$ is better known as \Box_{λ}^* .

Note that \Box_{λ} holds iff there exists a coherent λ -bounded *C*-sequence over λ^+ , and that \Box^*_{λ} holds iff there exists a weakly coherent λ -bounded *C*-sequence over λ^+ . The following terminology is also quite useful.

Definition 3.7. A C-sequence $\vec{C} = \langle C_{\delta} | \delta < \lambda^+ \rangle$ is a transversal for \Box_{λ}^* iff it is λ -bounded and weakly coherent.

A special case of Fact 3.4 states that if $\Box(\kappa)$ holds and $\kappa \geq \aleph_2$, then for every stationary $S \subseteq \kappa$, there exists a $\Box(\kappa)$ -sequence \vec{C} such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}(S,\kappa)$.⁵ Replacing $\Box(\kappa)$ by \Box_{λ} , better forms of guessing are available:

Fact 3.8 ([Rin14c, Corollary 2.4]). Suppose that λ is an uncountable cardinal.

Then \Box_{λ} holds iff there exists a coherent λ -bounded C-sequence $\langle C_{\delta} | \delta < \lambda^+ \rangle$ with the feature that for every club $D \subseteq \lambda^+$ and every $\sigma \in \operatorname{acc}(\lambda)$, there exists some $\delta < \lambda^+$ with $\operatorname{otp}(C_{\delta}) = \sigma$ such that $C_{\delta} \subseteq D$.

Remark 3.9. Note that for a \vec{C} as above, the map $\delta \mapsto \operatorname{otp}(C_{\delta})$ yields a canonical partition of $\operatorname{acc}(\lambda^+)$ into λ -many pairwise disjoint stationary sets.

An inspection of the proofs of [Rin14c, Lemma 2.8] and Proposition 2.22 makes it clear that the following holds true.

Theorem 3.10. Suppose that λ is an uncountable cardinal, and S, T are stationary subsets of λ^+ . Suppose also that either $S \cap E_{\neq cf(\lambda)}^{\lambda^+}$ is stationary or that Tr(S) is stationary. Then:

- (1) \Box_{λ} holds iff there exists a coherent λ -bounded C-sequence $\langle C_{\delta} | \delta < \lambda^+ \rangle$ with the feature that for every club $D \subseteq \lambda^+$, there exists a $\delta \in S$ such that $\operatorname{nacc}(C_{\delta}) \subseteq D \cap T$;
- (2) \square_{λ}^{*} holds iff there exists a weakly coherent λ -bounded C-sequence $\langle C_{\delta} | \delta < \lambda^{+} \rangle$ with the feature that for every club $D \subseteq \lambda^{+}$, there exists a $\delta \in S$ such that $\operatorname{nacc}(C_{\delta}) \subseteq D \cap T$.

We now turn to present another postprocessing function.

Lemma 3.11 (see [BR21, Lemma 4.9]). For every function $f : \kappa \to [\kappa]^{<\omega}$, the operator $\Phi_f : \mathcal{K}(\kappa) \to \mathcal{K}(\kappa)$ defined via:

$$\Phi_f(x) := x \cup \bigcup \{ f(\gamma) \cap (\sup(x \cap \gamma), \gamma) \mid \gamma \in \operatorname{nacc}(x) \}$$

is a postprocessing function.

The definition of Φ_f is motivated by the *regressive functions ideal* $J[\kappa]$ from [Rin17], and the following extension of it from [Rin22].

16

⁵Replacing κ by an abstract stationary set T is trickier, but note that if S is a stationary subset of $E_{>\sigma}^{\kappa}$ that reflects stationarily often, then for every $\Box(\kappa)$ -sequence \vec{C} such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}(S, T, \sigma)$, it is also the case that $\vec{C} \upharpoonright \operatorname{Tr}(S)$ witnesses $\operatorname{CG}(\operatorname{Tr}(S), T, \sigma)$.

Definition 3.12 ([Rin22]). $J_{\omega}[\kappa]$ stands for the collection of all subsets $S \subseteq \kappa$ for which there exist a club $C \subseteq \kappa$ and a sequence of functions $\langle f_i : \kappa \to [\kappa]^{<\omega} | i < \kappa \rangle$ with the property that for every $\delta \in S \cap C$, every regressive function $f : \delta \to \delta$, and every cofinal subset $\Gamma \subseteq \delta$, there exists an $i < \delta$ such that

$$\sup\{\gamma \in \Gamma \mid f(\gamma) \in f_i(\gamma)\} = \delta.$$

The next lemma gives a sufficient condition for moving from $CG_{\xi}(S,\kappa)$ to $CG_{\xi}(S,T)$.

Lemma 3.13. Suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ is a *C*-sequence witnessing $\operatorname{CG}_{\xi}(S, \kappa)$. If $S \in J_{\omega}[\kappa]$, then for every stationary $T \subseteq \kappa$, there exists a function $f : \kappa \to [\kappa]^{<\omega}$ such that $\langle \Phi_f(C_{\delta}) \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, T)$

Proof. Without loss of generality, $S \subseteq \operatorname{acc}(\kappa)$. Suppose that $S \in J_{\omega}[\kappa]$, and fix a club $C \subseteq \kappa$ and a sequence of functions $\langle f_i : \kappa \to [\kappa]^{<\omega} \mid i < \kappa \rangle$ as in Definition 3.12. For all $i < \kappa$ and $\delta \in S$, denote $C^i_{\delta} := \Phi_{f_i}(C_{\delta})$, so that $\operatorname{otp}(C^i_{\delta}) \leq \operatorname{otp}(C_{\delta})$.

Let T be an arbitrary stationary subset of κ .

Claim 3.13.1. There is an $i < \kappa$ such that $\langle C_{\delta}^i | \delta \in S \rangle$ witnesses CG(S,T).

Proof. Suppose not. For each $i < \kappa$, pick a club $D_i \subseteq \kappa$ such that for every $\delta \in S$,

$$\sup(\operatorname{nacc}(C^i_{\delta}) \cap D_i \cap T) < \delta.$$

Consider the two clubs $D := C \cap \Delta_{i < \kappa} D_i$ and $D' := \operatorname{acc}^+(D \cap T)$. By the choice of \vec{C} , pick $\delta \in S$ such that $\Gamma := \operatorname{nacc}(C_{\delta}) \cap D'$ is cofinal in δ . As Γ is a subset of $\operatorname{nacc}(C_{\delta}) \cap D'$, we may define a regressive function $f : \Gamma \to \delta$ via:

$$f(\gamma) := \min\{\beta \in D \cap T \mid \sup(C_{\delta} \cap \gamma) < \beta < \gamma\}.$$

As $\gamma \in S \cap C$, find $i < \delta$ such that $\Gamma' := \{\gamma \in \Gamma \mid f(\gamma) \in f_i(\gamma)\}$ is cofinal in δ . By possibly omitting an initial segment of Γ' , we may assume that $\sup(C_{\delta} \cap \min(\Gamma')) >$ i. Recalling the definition of D, it follows that for every $\gamma \in \Gamma'$, $f(\gamma) \in D_i \cap T$. So, for every $\gamma \in \Gamma'$, if we let $\beta := \sup(C_{\delta} \cap \gamma)$, then $C_{\delta}^i \cap (\beta, \gamma)$ is equal to the finite set $f_i(\gamma) \cap (\beta, \gamma)$ that contains $f(\gamma)$ which is an element of $D_i \cap T$. So $\sup(\operatorname{nacc}(C_{\delta}^i) \cap D_i \cap T) = \delta$, contradicting the choice of D_i .

Let *i* be given by the preceding claim. Then $f := f_i$ is as sought.

By [Rin22, Proposition 3.3], $J_{\omega}[\lambda^+]$ contains no stationary subsets of $E_{cf(\lambda)}^{\lambda^+}$. In particular, $J_{\omega}[\omega_1]$ is empty. So, unlike Fact 3.4, in the following we don't need to explicitly require κ to be $\geq \aleph_2$.

Corollary 3.14. If $\Box(\kappa)$ holds, then for every stationary $S \in J_{\omega}[\kappa]$ and every stationary $T \subseteq \kappa$, there exists a $\Box(\kappa)$ -sequence \vec{C} such that $\vec{C} \upharpoonright S$ witnesses CG(S,T).

We have described a way for moving from $CG(S, \kappa)$ to CG(S, T). Our next goal is to describe a way for moving from $CG(\kappa, T)$ to CG(S, T). First, a definition.

Definition 3.15. We say that a *C*-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ is \sqsubseteq^* -coherent iff for all $\delta \in S$ and $\bar{\delta} \in \operatorname{acc}(C_{\delta})$, it is the case that $\bar{\delta} \in S$ and $\sup(C_{\bar{\delta}} \triangle (C_{\delta} \cap \bar{\delta})) < \bar{\delta}$.

Remark 3.16. Every \sqsubseteq^* -coherent C-sequence is weakly coherent.

To preserve \sqsubseteq^* -coherence, one needs to consider a strengthening of Definition 3.1.

Definition 3.17. An operator $\Phi : \mathcal{K}(\kappa) \to \mathcal{K}(\kappa)$ is a *postprocessing*^{*} function if for every $x \in \mathcal{K}(\kappa)$, Clause (i)–(iii) of Definition 3.1 hold true, and, in addition:

(iv) for every $\bar{x} \in \mathcal{K}(\kappa)$ such that $\bar{x} \sqsubseteq^* x$ and $\sup(\bar{x}) \in \operatorname{acc}(\Phi(x)), \Phi(\bar{x}) \sqsubseteq^* \Phi(x)$.

It is readily checked that all the postprocessing functions we have met so far are moreover postprocessing^{*} functions. An example of a postprocessing function that is not postprocessing^{*} function may be found in [LHR19, Lemma 3.8]. In fact, it is unknown at present how to obtain the same effect of that map using a postprocessing^{*} function.

Theorem 3.18. Suppose that $\kappa \geq \aleph_2$, and that there exists an \sqsubseteq^* -coherent *C*-sequence witnessing $CG(\kappa, T)$.

For every stationary $S \subseteq \kappa$, there exists an \sqsubseteq^* -coherent C-sequence over κ whose restriction to S witnesses CG(S,T).

Proof. The proof will be an adaptation of the proof of [Rin17, Theorem 4.13]. Suppose towards a contradiction that S is a counterexample. Fix an \sqsubseteq^* -coherent C-sequence $\vec{e} = \langle e_{\delta} \mid \delta < \kappa \rangle$ witnessing $\operatorname{CG}(\kappa, T)$. By recursion on $i < \omega_1$, we construct a club $D_i \subseteq \kappa$ and an \sqsubseteq^* -coherent C-sequence $\vec{C}^i = \langle C^i_{\alpha} \mid \alpha < \kappa \rangle$. Our construction will have the property that for all $\alpha < \kappa$ and $j < i < \omega_1, C^i_{\alpha} \subseteq C^i_{\alpha}$.

For i = 0, let $D_0 := \kappa$ and $\vec{C^0} := \vec{e}$.

► For every $i < \omega_1$ such that a club $D_i \subseteq \kappa$ and an \sqsubseteq^* -coherent *C*-sequence \vec{C}^i have been constructed, by the assumption we can find a club F^{D_i} such that, for every $\delta \in S$,

$$\sup(\operatorname{nacc}(C^i_{\delta}) \cap F^{D_i} \cap T) < \delta.$$

So let $D_{i+1} := D_i \cap F^{D_i}$. As for constructing C^{i+1} , we do this by recursion on $\alpha < \kappa$. To start, let $C_0^{i+1} := \emptyset$, and for every $\alpha < \kappa$ let $C_{\alpha+1}^{i+1} := \{\alpha\}$. Next, for $\alpha \in \operatorname{acc}(\kappa)$,

$$C^{i+1}_{\alpha} := C^i_{\alpha} \cup \bigcup \{ C^{i+1}_{\gamma} \setminus \sup(C^i_{\alpha} \cap \gamma) \mid \gamma \in \operatorname{nacc}(C^i_{\alpha}) \setminus (F^{D_i} \cap T) \}.$$

It is clear that $C^{i+1} = \langle C^{i+1}_{\alpha} | \alpha < \kappa \rangle$ is \sqsubseteq^* -coherent as well.

► For every $i \in \operatorname{acc}(\omega_1)$ such that $\langle (D_j, \vec{C^j}) \mid j < i \rangle$ has been constructed as required, let $D_i := \bigcap_{j < i} D_j$ and for every $\alpha < \kappa$, let $C^i_\alpha := \bigcup_{j < i} C^j_\alpha$.

Claim 3.18.1. Let $\alpha < \kappa$. Then $\operatorname{acc}^+(C^i_\alpha) = \bigcup_{j < i} \operatorname{acc}(C^j_\alpha)$.

However, this implies that $\beta \notin \operatorname{acc}(C^i_{\alpha})$, which is a contradiction.

Proof. Let $\beta \in \operatorname{acc}^+(C^i_\alpha)$. The sequence $\langle \min(C^j_\alpha \setminus \beta) \mid j < i \rangle$ is weakly decreasing, and as i is a nonzero limit ordinal, it stabilizes at some $j^* < i$. Let β^+ be this stable value (which may be equal to β). If there exists $j \in [j^*, i)$ such that $\beta^+ \in \operatorname{acc}(C^j_\alpha)$, then in fact $\beta^+ = \beta$ and we can finish. So, suppose that this is not so. If there exists $j \in [j^*, i)$ such that $\beta^+ \notin F^{D_j} \cap T$, then $\beta^+ \in \operatorname{acc}(C^{j+1}_\alpha)$ and again we can finish. So suppose this is not so. Let $\beta^- := \sup(C^{j^*}_\alpha \cap \beta^+)$ so that $\beta^- < \beta \leq \beta^+$. Now examining the construction of C^j_α for $j \in (j^*, i)$, it is clear that for every $j \in [j^*, i)$,

$$(\beta^-,\beta^+] \cap C^{j^*}_{\alpha} = (\beta^-,\beta^+] \cap C^j_{\alpha}.$$

 \dashv

So $\vec{C^i} = \langle C^i_{\alpha} \mid \alpha < \kappa \rangle$ is a *C*-sequence. Furthermore, since, $\vec{C^j}$ is \sqsubseteq^* -coherent for every j < i, the preceding claim altogether implies that $\vec{C^i}$ is \sqsubseteq^* -coherent.

Consider the club $D := \bigcap_{i < \omega_1} D_i$. As \vec{C} witnesses $CG(\kappa, T)$, the following set is stationary:

$$B := \{\beta < \kappa \mid \sup(\operatorname{nacc}(C_{\beta}) \cap D \cap T) = \beta\}.$$

Notice that by the nature of our recursive construction, for all $i < \omega_1$,

 $\operatorname{nacc}(C_{\beta}) \cap D \cap T) \subseteq \operatorname{nacc}(C^{i}_{\beta}) \cap D \cap T,$

and hence, for all $\beta \in B$ and $i < \omega_1$,

$$\sup(\operatorname{nacc}(C^i_\beta) \cap D \cap T) = \beta.$$

Pick $\delta \in S \cap \operatorname{acc}^+(B)$. For each $i < \omega_1$ the following ordinal is smaller than δ

$$\epsilon_i := \sup(\operatorname{nacc}(C^i_{\delta}) \cap F^{D_i} \cap T).$$

We now perform a case analysis to reach a contradiction.

CASE 1. $cf(\delta) > \aleph_0$. Let $\epsilon^* := \sup_{i < \omega} \epsilon_i$, so that $\epsilon^* < \delta$. Pick $\beta \in (\epsilon^*, \delta) \cap B$. For every $i < \omega$, let $\gamma_i := \min(C_{\delta}^i \setminus \beta)$ so that $\langle \gamma_i \mid i < \omega \rangle$ is weakly decreasing. Then pick $i < \omega$ such that $\gamma_i = \gamma_{i+1}$.

SUBCASE 1.1. $\gamma_i > \beta$. It follows that $\gamma_i \in \operatorname{nacc}(C^i_{\delta})$. As $\epsilon^* < \beta < \gamma_i$, we have that $\gamma_i \notin F^{D_i} \cap T$. It follows from our recursive construction that $C^{i+1}_{\delta} \cap [\beta, \gamma_i) = C^{i+1}_{\gamma_i} \cap [\beta, \gamma_i)$ and the latter set is nonempty so that $\gamma_{i+1} < \gamma_i$ which is a contradiction.

SUBCASE 1.2. $\gamma_i = \beta$ and $\beta \in \text{nacc}(C^i_{\delta})$. So γ_i is an element of $\text{nacc}(C^i_{\delta})$ above ϵ^* , and we are back to Subcase 1.1.

SUBCASE 1.3. $\gamma_i = \beta$ and $\beta \in \operatorname{acc}(C^i_{\delta})$. In this case, by \sqsubseteq^* -coherence, we have that $\sup((C^i_{\beta} \bigtriangleup C^i_{\delta}) \cap \beta) < \beta$. This implies that

$$\sup(\operatorname{nacc}(C^i_{\delta}) \cap \beta \cap D \cap T) = \beta.$$

But $D \subseteq F^{D_i}$, contradicting the fact that $\epsilon_i < \beta$.

SUBCASE 2. $cf(\delta) = \aleph_0$. Find an uncountable $I \subseteq \omega_1$ such that

 $\epsilon^* := \sup\{\max\{\epsilon_i, \epsilon_{i+1}\} \mid i \in I\}$

is smaller than δ . Pick $\beta \in (\epsilon^*, \delta) \cap B$. For every $i < \omega_1$, let $\gamma_i := \min(C^i_{\delta} \setminus \beta)$ so that $\langle \gamma_i \mid i < \omega_1 \rangle$ is weakly decreasing. Pick a large enough $i \in I$ such that $\gamma_i = \gamma_{i+1}$.

SUBCASE 2.1. $\gamma_i > \beta$. Same as in Subcase 1.1.

SUBCASE 2.2. $\gamma_i = \beta$ and $\beta \in \operatorname{nacc}(C^i_{\delta})$. Same as in Subcase 1.2. **SUBCASE 2.3.** $\gamma_i = \beta$ and $\beta \in \operatorname{acc}(C^i_{\delta})$. Same as in Subcase 1.3.

Corollary 3.19. If $\Box(\kappa)$ holds and $J_{\omega}[\kappa]$ contains a stationary set, then for all stationary subsets S, T of κ , there exists an \sqsubseteq^* -coherent C-sequence $\vec{C} = \langle C_{\delta} | \delta < \kappa \rangle$ such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}(S, T)$.

Recalling Theorem 3.10, we now turn to deal with stationary subsets of $E_{cf(\lambda)}^{\lambda^+}$, dividing the results into two, depending on whether λ is regular or singular.

Theorem 3.20. Suppose that λ is a regular uncountable cardinal, and \Box_{λ}^* holds.

For every stationary $S \subseteq E_{\lambda}^{\lambda^+}$, there exists a transversal $\vec{C} = \langle C_{\delta} \mid \delta < \lambda^+ \rangle$ for \Box_{λ}^* such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}_{\lambda}(S, E_{\lambda}^{\lambda^+})$.

Proof. Suppose not, and fix a stationary $S \subseteq E_{\lambda}^{\lambda^+}$ that constitutes a counterexample. As \Box_{λ}^* holds, we may fix a transversal $\vec{e} = \langle e_{\delta} | \delta < \lambda^+ \rangle$ for \Box_{λ}^* . We shall recursively construct a sequence $\langle (D_n, \vec{e^n}, \vec{C^n}) \mid n < \omega \rangle$ such that D_n is a club in λ^+ , and $\vec{e^n}$ and $\vec{C^n}$ are transversals for \Box^*_{λ} .

Set $D_0 := \operatorname{acc}(\lambda^+)$, $\vec{e^0} := \vec{e}$ and $\vec{C^0} := \vec{e}$. Next, suppose that $n < \omega$ and that $\langle (D_j, \vec{e^j}, \vec{C^j}) \mid j \leq n \rangle$ has already been successfully defined. As $\vec{C^n} = \langle C^n_{\delta} \mid \delta < \lambda^+ \rangle$ is a transversal for \Box_{λ}^{*} , by the choice of the stationary set S, it follows that we may pick a subclub $D_{n+1} \subseteq D_n$ such that, for every $\delta \in S$,

$$\sup(\operatorname{nacc}(C^n_{\delta}) \cap D_{n+1} \cap E^{\lambda^+}_{\lambda}) < \delta.$$

Consider the postprocessing function $\Phi_{D_{n+1}}$ from Definition 2.12. For every $\delta \in \lambda^+ \setminus S$, let $e_{\delta}^{n+1} := e_{\delta}$ and $C_{\delta}^{n+1} := e_{\delta}$. For every $\delta \in S$, let $e_{\delta}^{n+1} := \Phi_{D_{n+1}}(C_{\delta}^n)$, and then let

$$C_{\delta}^{n+1} := e_{\delta}^{n+1} \cup \{e_{\gamma} \setminus \sup(e_{\delta}^{n+1} \cap \gamma) \mid \gamma \in \operatorname{nacc}(e_{\delta}^{n+1}) \cap E_{<\delta}^{\lambda^{+}}\}.$$

By [BR19b, Lemma 2.8], $e^{\vec{n+1}} := \langle e^{n+1}_{\delta} | \delta < \lambda^+ \rangle$ is again a transversal for \Box^*_{λ} .

Claim 3.20.1. $\vec{C^{n+1}}$ is a transversal for \Box_{λ}^* .

Proof. Since $e^{\vec{n+1}}$ is a λ -bounded C-sequence, the definition of $C^{\vec{n+1}}$ makes it clear that it is as well. Suppose that $C^{\vec{n}+1}$ is not weakly coherent, and pick the least $\alpha < \lambda^+$ such that

$$|\{C_{\delta}^{n+1} \cap \alpha \mid \delta < \lambda^+\}| = \lambda^+.$$

As $C^{\vec{n}+1} \upharpoonright (\lambda^+ \setminus S) = e^{\vec{n}+1} \upharpoonright (\lambda^+ \setminus S)$, and $e^{\vec{n}+1}$ is a transversal for \Box^*_{λ} , it follows that

$$|\{e_{\delta}^{n+1} \cap \alpha \mid \delta \in S\}| < |\{C_{\delta}^{n+1} \cap \alpha \mid \delta \in S\}| = \lambda^{+},$$

so we may fix $\Delta \in [S]^{\lambda^+}$ such that:

- δ → Cⁿ⁺¹_δ ∩ α is injective over Δ, but
 δ → eⁿ⁺¹_δ ∩ α is constant over Δ.

Fix $\epsilon < \alpha$ such that $\sup(e_{\delta}^{n+1} \cap \alpha) = \epsilon$ for all $\delta \in \Delta$. By minimality of α , and by possibly shrinking Δ further, we may also assume that

• $\delta \mapsto C_{\delta}^{n+1} \cap \epsilon$ is constant over Δ .

It thus follows from the definition of C^{n+1} that the map $\delta \mapsto C^{n+1}_{\delta} \cap [\epsilon, \alpha)$ is injective over Δ , and that, for every $\delta \in \Delta$, $C^{n+1}_{\delta} \cap [\epsilon, \alpha) = e_{\gamma} \cap [\epsilon, \alpha)$ for $\gamma :=$ $\min(e_s^{n+1} \setminus (\epsilon+1))$. In particular,

$$|\{e_{\gamma} \cap [\epsilon, \alpha) \mid \gamma < \lambda^{+}\}| = \lambda^{+}$$

contradicting the fact that \vec{e} is a transversal for a \Box_{λ}^* -sequence.

 \dashv

This completes the construction of the sequence $\langle (D_n, \vec{e^n}, \vec{C^n}) \mid n < \omega \rangle$. Now, let $D := \bigcap_{n < \omega} D_n$. Pick $\delta \in S$ such that $\operatorname{otp}(D \cap \delta) = \omega^{\delta} > \lambda$. Recall that, for every $n < \omega$, the following ordinal is smaller than δ :

$$\epsilon_n := \sup(\operatorname{nacc}(C^n_\delta) \cap D_{n+1} \cap E^{\lambda^+}_\lambda).$$

Since $\operatorname{cf}(\lambda) > \omega$, for every $\alpha < \delta$, $\operatorname{otp}(\bigcup_{n < \omega} C_{\delta}^n \cap \alpha) < \lambda$. So, $\operatorname{otp}(\bigcup_{n < \omega} C_{\delta}^n) = 0$ $\lambda < \omega^{\delta} = \operatorname{otp}(D \cap \delta)$, and we may fix $\beta \in D \setminus \bigcup_{n < \omega} C_{\delta}^{n}$ above $\sup_{n < \omega} \epsilon_{n}$. Clearly, for each $n < \omega$, $\gamma_n := \min(C^n_{\delta} \setminus \beta)$ is an element of $\operatorname{nacc}(C^n_{\delta})$ above β .

Let $n < \omega$. Since $e_{\delta}^{n+1} = \Phi_{D_{n+1}}(C_{\delta}^n)$ and $\sup(D_{n+1} \cap \delta) = \delta$,

$$e_{\delta}^{n+1} = \{ \sup(D_{n+1} \cap \eta) \mid \eta \in C_{\delta}^{n}, \eta > \min(D_{n+1}) \}.$$

In particular, $\sup(D_{n+1} \cap \gamma_n) \in e_{\delta}^{n+1}$. As $\gamma_n > \beta$ and $\beta \in D \subseteq D_{n+1}$, it is the case that $\beta \leq \sup(D_{n+1} \cap \gamma_n)$. As $e_{\delta}^{n+1} \subseteq C_{\delta}^{n+1}$, altogether,

$$\beta < \gamma_{n+1} = \min(C_{\delta}^{n+1} \setminus \beta) \le \min(e_{\delta}^{n+1} \setminus \beta) = \sup(D_{n+1} \cap \gamma_n) \le \gamma_n.$$

Now, pick $n < \omega$ such that $\gamma_{n+1} = \gamma_n$. There are two options, each leads to a contradiction.

► If $\gamma_{n+1} \in e_{\delta}^{n+1}$, then since $e_{\delta}^{n+1} \subseteq C_{\delta}^{n+1}$, and $\gamma_{n+1} \in \operatorname{nacc}(C_{\delta}^{n+1})$, $\gamma_{n+1} \in \operatorname{nacc}(e_{\delta}^{n+1})$. As, $\gamma_{n+1} \in e_{\delta}^{n+1} \subseteq D_{n+1} \subseteq D_0$, γ is a limit ordinal. Since $C_{\delta}^{n+1} \cap [\beta, \gamma_{n+1}]$ is empty, the definition of C_{δ}^{n+1} implies that $\operatorname{cf}(\gamma_{n+1}) = \lambda$. Altogether, $\gamma_{n+1} \in \operatorname{nacc}(C_{\delta}^{n+1}) \cap D_{n+1} \cap E_{\delta}^{\lambda^+}$, contradicting the fact that $\gamma_{n+1} \geq \beta \geq \epsilon_{n+1}$.

 $\gamma_{n+1} \in \operatorname{nacc}(C_{\delta}^{n+1}) \cap D_{n+1} \cap E_{\lambda}^{\lambda^{+}}$, contradicting the fact that $\gamma_{n+1} > \beta > \epsilon_{n+1}$. \blacktriangleright If $\gamma_{n+1} \notin e_{\delta}^{n+1}$, then since $\gamma_{n+1} = \gamma_n \in C_{\delta}^n$, the definition of e_{δ}^{n+1} implies that $\gamma_{n+1} < \sup(D_{n+1} \cap \gamma_n)$. So, this time,

$$\gamma_{n+1} = \min(C_{\delta}^{n+1} \setminus \beta) < \min(e_{\delta}^{n+1} \setminus \beta) = \sup(D_{n+1} \cap \gamma_n) \le \gamma_n,$$

contradicting the choice of n.

In order to obtain a correct analogue of the preceding result, we introduce the following natural strengthening of Definition 2.2, in which we replace the stationary set $T \subseteq \kappa$ by a sequence $\vec{T} = \langle T_i \mid i < \theta \rangle$ of stationary subsets of κ .

Definition 3.21. $\operatorname{CG}_{\xi}(S, \vec{T}, \sigma, \vec{J})$ asserts the existence of a ξ -bounded *C*-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ such that, for every club $D \subseteq \kappa$ there is a $\delta \in S$ such that for every $i < \min\{\delta, \theta\}$,

$$\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T_i\} \in J_{\delta}^+.$$

Convention 3.22. Convention 2.3 applies to the above definition, as well.

Theorem 3.23. Suppose that λ is a singular cardinal of uncountable cofinality, and \Box_{λ}^* holds. Let $\langle \lambda_i | i < cf(\lambda) \rangle$ be the increasing enumeration of a club in λ .

For every stationary $S \subseteq E_{\geq \omega}^{\lambda^+}$, there exists a transversal $\vec{C} = \langle C_{\delta} \mid \delta < \lambda^+ \rangle$ for \Box_{λ}^* such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}_{\lambda}(S, \langle E_{\geq \lambda_i}^{\lambda^+} \mid i < \operatorname{cf}(\lambda) \rangle)$.

Proof. Suppose not, and fix a stationary $S \subseteq E_{>\omega}^{\lambda^+}$ that constitutes a counterexample. Without loss of generality, $\min(S) \ge \lambda$. As \Box_{λ}^{*} holds, we may fix a transversal $\vec{e} = \langle e_{\delta} \mid \delta < \lambda^+ \rangle$ for \Box_{λ}^{*} . As λ is singular, we may assume that $\operatorname{otp}(e_{\delta}) < \lambda$ for every $\delta < \lambda^+$ (e.g., by appealing to [BR19a, Lemma 3.1] with $\Sigma := \{\lambda_i \mid i < \operatorname{cf}(\lambda)\}$).

Following the proof approach of Theorem 3.20, we shall recursively construct a sequence $\langle (D_n, \vec{e^n}, \vec{C^n}) | n < \omega \rangle$ such that D_n is a club in λ^+ , and $\vec{e^n}$ and $\vec{C^n}$ are transversals for \Box^*_{λ} .

Set $D_0 := \operatorname{acc}(\lambda^+)$, $\vec{e^0} := \vec{e}$ and $\vec{C^0} := \vec{e}$. Next, suppose that $n < \omega$ and that $\langle (D_j, \vec{e^j}, \vec{C^j}) \mid j \leq n \rangle$ has already been successfully defined. As $\vec{C^n} = \langle C^n_{\delta} \mid \delta < \lambda^+ \rangle$ is a transversal for \Box^*_{λ} , by the choice of the stationary set S, it follows that we may pick a subclub $D_{n+1} \subseteq D_n$ such that, for every $\delta \in S$, for some $i^{n+1}_{\delta} < \operatorname{cf}(\lambda)$,

$$\sup(\operatorname{nacc}(C^n_{\delta}) \cap D_{n+1} \cap E^{\lambda^+}_{\geq \lambda_{i^{n+1}_{\delta}}}) < \delta.$$

Consider the postprocessing function $\Phi_{D_{n+1}}$ from Definition 2.12. For every $\delta \in \lambda^+ \setminus S$, let $e_{\delta}^{n+1} := e_{\delta}$ and $C_{\delta}^{n+1} := e_{\delta}$. For every $\delta \in S$, let $e_{\delta}^{n+1} := \Phi_{D_{n+1}}(C_{\delta}^n)$, and then let

$$C^{n+1}_{\delta} := e^{n+1}_{\delta} \cup \{e_{\gamma} \setminus \sup(e^{n+1}_{\delta} \cap \gamma) \mid \gamma \in \operatorname{nacc}(e^{n+1}_{\delta}) \cap E^{\lambda^{+}}_{<\lambda_{i^{n+1}_{\delta}}} \}.$$

By [BR19b, Lemma 2.8], $e^{\vec{n}+1} := \langle e^{n+1}_{\delta} | \delta < \lambda^+ \rangle$ is again a transversal for \Box^*_{λ} . By the exactly same proof of Claim 3.20.1, also $C^{\vec{n}+1} := \langle C^{n+1}_{\delta} | \delta < \lambda^+ \rangle$ is a transversal for \Box^*_{λ} . Furthermore, $\operatorname{otp}(e^{n+1}_{\delta}) \leq \operatorname{otp}(C^{n+1}_{\delta}) < \lambda$ for all $\delta < \lambda^+$.

This completes the construction of the sequence $\langle (D_n, \vec{e^n}, \vec{C^n}) | n < \omega \rangle$. Now, let $D := \bigcap_{n < \omega} D_n$. Pick $\delta \in S$ such that $\operatorname{otp}(D \cap \delta) = \omega^{\delta} > \lambda$. Recall that, for every $n < \omega$, the following ordinal is smaller than δ :

$$\epsilon_n := \sup(\operatorname{nacc}(C^n_\delta) \cap D_{n+1} \cap E^{\lambda^+}_{\geq \lambda_{i_s^{n+1}}}).$$

As $\operatorname{cf}(\lambda) > \omega$, $\operatorname{otp}(\bigcup_{n < \omega} C_{\delta}^n) < \lambda$, so, we may fix $\beta \in D \setminus \bigcup_{n < \omega} C_{\delta}^n$ above $\sup_{n < \omega} \epsilon_n$. Clearly, for each $n < \omega$, $\gamma_n := \min(C_{\delta}^n \setminus \beta)$ is an element of $\operatorname{nacc}(C_{\delta}^n)$ above β .

Let $n < \omega$. Since $e_{\delta}^{n+1} = \Phi_{D_{n+1}}(C_{\delta}^n)$ and $\sup(D_{n+1} \cap \delta) = \delta$,

$$e_{\delta}^{n+1} = \{ \sup(D_{n+1} \cap \eta) \mid \eta \in C_{\delta}^n, \eta > \min(D_{n+1}) \}.$$

In particular, $\sup(D_{n+1} \cap \gamma_n) \in e_{\delta}^{n+1}$. As $\gamma_n > \beta$ and $\beta \in D \subseteq D_{n+1}$, it is the case that $\beta \leq \sup(D_{n+1} \cap \gamma_n)$. As $e_{\delta}^{n+1} \subseteq C_{\delta}^{n+1}$, altogether,

$$\beta < \gamma_{n+1} = \min(C_{\delta}^{n+1} \setminus \beta) \le \min(e_{\delta}^{n+1} \setminus \beta) = \sup(D_{n+1} \cap \gamma_n) \le \gamma_n.$$

Now, pick $n < \omega$ such that $\gamma_{n+1} = \gamma_n$. There are two options, each leads to a contradiction.

► If $\gamma_{n+1} \in e_{\delta}^{n+1}$, then since $e_{\delta}^{n+1} \subseteq C_{\delta}^{n+1}$, and $\gamma_{n+1} \in \operatorname{nacc}(C_{\delta}^{n+1})$, $\gamma_{n+1} \in \operatorname{nacc}(e_{\delta}^{n+1})$. As, $\gamma_{n+1} \in e_{\delta}^{n+1} \subseteq D_{n+1} \subseteq D_0$, γ is a limit ordinal. So, since $C_{\delta}^{n+1} \cap [\beta, \gamma_{n+1})$ is empty, the definition of C_{δ}^{n+1} implies that $\operatorname{cf}(\gamma_{n+1}) \ge \lambda_{i_{\delta}^{n+1}}$. Altogether, $\gamma_{n+1} \in \operatorname{nacc}(C_{\delta}^{n+1}) \cap D_{n+1} \cap E_{\lambda_{i_{\delta}^{n+1}}}^{\lambda^+}$, contradicting the fact that $\gamma_{n+1} > \beta > \epsilon_{n+1}$.

• If $\gamma_{n+1} \notin e_{\delta}^{n+1}$, then since $\gamma_{n+1} = \gamma_n \in C_{\delta}^n$, the definition of e_{δ}^{n+1} implies that $\gamma_{n+1} < \sup(D_{n+1} \cap \gamma_n)$. So, this time,

$$\gamma_{n+1} = \min(C_{\delta}^{n+1} \setminus \beta) < \min(e_{\delta}^{n+1} \setminus \beta) = \sup(D_{n+1} \cap \gamma_n) \le \gamma_n,$$

contradicting the choice of n.

By applying the proof of Proposition 2.22 on the *C*-sequence produced by the preceding, we get a somewhat cleaner form of guessing, as follows.

Corollary 3.24. Suppose that λ is a singular cardinal of uncountable cofinality, and \Box^*_{λ} holds. For every stationary $S \subseteq E^{\lambda^+}_{>\omega}$, there exists a transversal $\vec{C} = \langle C_{\delta} | \delta < \lambda^+ \rangle$ for \Box^*_{λ} satisfying the following:

- for every $\delta < \lambda^+$, $\operatorname{otp}(C_{\delta}) < \lambda$;
- for every club $D \subseteq \lambda^+$, there exists $\delta \in S$ such that $C_{\delta} \subseteq D$ and $\sup(\operatorname{nacc}(C_{\delta}) \cap D \cap E_{>\mu}^{\lambda^+}) = \delta$ for every $\mu < \lambda$.

3.1. When coherence is not available. By waiving any coherence considerations, the proofs of Theorems 3.20 and 3.23 (together with Proposition 2.22) yield, respectively, the general case of the introduction's Fact 1.4 and a result from [ES09].

Fact 3.25 ([She94c, Claim 2.4]). For every regular uncountable cardinal λ , for every stationary $S \subseteq E_{\lambda}^{\lambda^+}$, $\operatorname{CG}_{\lambda}(S, E_{\lambda}^{\lambda^+})$ holds. Furthermore, for every triple $\lambda \leq \nu < \kappa$ of regular uncountable cardinals, for every stationary $S \subseteq E_{\lambda}^{\kappa}$, $\operatorname{CG}_{\nu}(S, E_{\geq \nu}^{\kappa})$ holds.

Fact 3.26 ([ES09, Theorem 2]). For every singular cardinal λ of uncountable cofinality and every stationary $S \subseteq E_{cf(\lambda)}^{\lambda^+}$, there exists a $cf(\lambda)$ -bounded C-sequence $\langle C_{\delta} | \delta \in S \rangle$ satisfying the following.

For every club $D \subseteq \lambda^+$, there exists $\delta \in S$ such that $C_{\delta} \subseteq D$ and $\langle cf(\gamma) | \gamma \in nacc(C_{\delta}) \rangle$ is strictly increasing and converging to λ .

Likewise, by changing the choice of the initial C-sequence \vec{e} in the proof of Theorem 3.18, one obtains a proof of the following.

Theorem 3.27 (Shelah). Suppose that R, S, T are stationary subsets of a regular cardinal $\kappa \geq \aleph_2$.

- (1) If T is a nonreflecting stationary set, then CG(S,T) holds;
- (2) If R is a nonreflecting stationary subset of $E_{\geq \sigma}^{\kappa}$, then $CG(R, T, \sigma)$ implies $CG(S, T, \sigma)$.

Remark 3.28. Note that even if $S \subseteq E_{\xi}^{\kappa}$, we still get $\operatorname{CG}(S,T)$, but not $\operatorname{CG}_{\xi}(S,T)$. Indeed, by the preceding corollary, if there exists a nonreflecting stationary subset of $E_{\aleph_0}^{\aleph_2}$, then $\operatorname{CG}(E_{\aleph_1}^{\aleph_2}, E_{\aleph_0}^{\aleph_2})$ holds. In contrast, running the forcing from [Asp14, Theorem 1.6] over a model of \Box_{ω_1} , one gets a generic extension with a nonreflecting stationary subset of $E_{\aleph_0}^{\aleph_2}$ in which $\operatorname{CG}_{\omega_1}(E_{\aleph_1}^{\aleph_2}, E_{\aleph_0}^{\aleph_2})$ fails.

Corollary 3.29. Suppose $\sigma < \sigma^+ < \kappa$ are infinite regular cardinals, and E^{κ}_{σ} admits a nonreflecting stationary set. For every stationary $S \subseteq \kappa$:

- (1) $CG(S, \kappa, \sigma)$ holds;
- (2) If $\kappa = \lambda^+$ and $\sigma \neq cf(\lambda)$, then $CG(S, T, \sigma)$ holds for every stationary $T \subseteq \kappa$.

Proof. Let R be a nonreflecting stationary subset of E_{σ}^{κ} , By Theorem 2.14, $CG(R, \kappa, \sigma)$ holds. By Corollary 2.23, if $\kappa = \lambda^+$ and $\sigma \neq cf(\lambda)$, then furthermore $CG(R, T, \sigma)$ holds for every stationary $T \subseteq \kappa$. Now appeal to Corollary 3.27(2).

Forgetting about coherence, Corollary 3.19 has the following strong consequence.

Corollary 3.30. Suppose that $\Box(\lambda^+)$ holds, and one of the following:

•
$$\lambda > \exists \dots$$

•
$$\lambda^{\aleph_0} = \lambda$$
:

- $\lambda = \mathfrak{b} = \aleph_1$;
- $\lambda \geq 2^{\aleph_1}$ and Shelah's Strong Hypothesis (SSH) holds;
- There exists an infinite regular cardinal θ such that $2^{\theta} \leq \lambda < \theta^{+\theta}$.

Then CG(S,T) holds for all stationary subsets S,T of λ^+ .

Proof. By Corollary 5.1, Corollary 5.3 and Corollary 5.7 of [Rin22], any of the above hypotheses imply that $J_{\omega}[\lambda^+]$ contains a stationary set.

TANMAY INAMDAR AND ASSAF RINOT

4. Partitioned the club-guessing

The theme of this section is partitioned club guessing as in Fact 1.7. The main definition is Definition 4.1, where various types of partitions are considered. The bulk of our results, with the exception of Subsection 4.2, however are about the stronger partitioning club-guessing. The difference being that in the former we obtain a C-sequence and a partition, whereas in the latter we are given the C-sequence in advance and then are given the task of partitioning it.

In Subsection 4.1, we partition club-guessing sequences using colouring principles from [IR22a, IR22b] which were in fact discovered while working on partitioning club-guessing. We show how these colouring principles allow for an abstract approach to partitioning club-guessing, separating the club-guessing technology from the combinatorial technology given to us by the relevant hypothesis.

In Subsection 4.2, we construct partitioned club-guessing using these colouring principles. Furthermore, we can obtain partitioned club-guessing sequences satisfying coherence features as well.

In Subsection 4.3, we list the results from [IR22a, IR22b] under which the colouring principles can be obtained, and draw conclusions. In particular, we find an higher analogue of a combinatorial construction on \aleph_1 due to Moore from [Moo08].

In Subsection 4.4, we address the problem of partitioning a club-guessing *C*-sequence \vec{C} over $S \subseteq \kappa$ into κ many guessing sequences $\langle \vec{C} \upharpoonright S_i \mid i < \kappa \rangle$.

Definition 4.1. For a *C*-sequence $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$, we define three sets of cardinals:

• $\Theta_0(\vec{C}, T, \sigma, \vec{J})$ denotes the set of all cardinals θ for which there exists a function $h: \kappa \to \theta$ satisfying the following.

For every club $D \subseteq \kappa$, there exists $\delta \in S$ such that, for every $\tau < \theta$,

 $\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \cap h^{-1}\{\tau\}\} \in J_{\delta}^+.$

• $\Theta_1(\vec{C}, T, \sigma, \vec{J})$ denotes the set of all cardinals θ for which there exists a function $h: \kappa \to \theta$ satisfying the following.

For every club $D \subseteq \kappa$, there exists $\delta \in S$ such that, for every $\tau < \theta$,

 $\{\beta < \delta \mid h(\operatorname{otp}(C_{\delta} \cap \beta)) = \tau \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} \in J_{\delta}^+.$

• $\Theta_2(\vec{C}, T, \sigma, \vec{J})$ denotes the set of all cardinals θ for which there exists a sequence of functions $\langle h_\delta : \delta \to \theta \mid \delta \in S \rangle$ satisfying the following. For every club $D \subseteq \kappa$, there exists $\delta \in S$ such that, for every $\tau < \theta$,

 $\{\beta < \delta \mid h_{\delta}(\beta) = \tau \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} \in J_{\delta}^+.$

Convention 4.2. Convention 2.3 applies to the above definition, as well.

Remark 4.3. $\Theta_0(\vec{C}, T, \sigma, \vec{J}) \subseteq \Theta_2(\vec{C}, T, \sigma, \vec{J})$ and $\Theta_1(\vec{C}, T, \sigma, \vec{J}) \subseteq \Theta_2(\vec{C}, T, \sigma, \vec{J})$.

Proposition 4.4. For a stationary $S \subseteq E_{\theta}^{\kappa}$ and a sequence \vec{C} witnessing $CG(S, T, \kappa)$, $\theta \in \Theta_2(\vec{C}, T, \sigma)$ for any choice of $\sigma < \theta$.

Lemma 4.5. Suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ is a *C*-sequence witnessing $\operatorname{CG}_{\xi}(S,T)$. For every $\theta \in \Theta_2(\vec{C},T)$ such that $\alpha + \beta < \xi$ for all $(\alpha,\beta) \in \theta \times \xi$, there exists a *C*-sequence $\vec{C}^{\bullet} = \langle C^{\bullet}_{\delta} \mid \delta \in S \rangle$ witnessing $\operatorname{CG}_{\xi}(S,T)$ for which $\theta \in \Theta_1(\vec{C}^{\bullet},T)$. *Proof.* Without loss of generality, $S \subseteq \operatorname{acc}(\kappa)$. Suppose that $\theta \in \Theta_2(\vec{C}, T)$ is such that $\alpha + \beta < \xi$ for all $(\alpha, \beta) \in \theta \times \xi$. Let $h : \kappa \to \theta$ be a surjection such that, for every $\epsilon < \kappa$, $\{h(\iota + 1) \mid \epsilon < \iota < \epsilon + \theta\} = \theta$. If θ is uncountable, we also require that $\{h(\omega \cdot \iota) \mid \epsilon < \iota < \epsilon + \theta\} = \theta$ for every $\epsilon < \kappa$. It follows that for each $(\gamma, \epsilon, \tau) \in \kappa \times \kappa \times \theta$, we may fix $y_{\gamma,\epsilon,\tau}$ such that:

- $y_{\gamma,\epsilon,\tau}$ is a closed nonempty subset of $E_{<\theta}^{\kappa}$;
- $\min(y_{\gamma,\epsilon,\tau}) = \gamma;$
- $\max(y_{\gamma,\epsilon,\tau}) < \gamma + \theta;$
- $h(\epsilon + \operatorname{otp}(y_{\gamma,\epsilon,\tau}) 1) = \tau.$

Fix a sequence $\langle h_{\delta} : \delta \to \theta \mid \delta \in S \rangle$ witnessing that $\theta \in \Theta_2(\vec{C}, T)$. We now construct $\vec{C}^{\bullet} = \langle C^{\bullet}_{\delta} \mid \delta \in S \rangle$ as follows. Given $\delta \in S$, let $\langle \delta_i \mid i < \operatorname{otp}(C_{\delta}) \rangle$ denote the increasing enumeration of $\{0\} \cup C_{\delta}$. Construct a sequence $\langle x_i^{\delta} \mid i < \operatorname{otp}(C_{\delta}) \rangle$ by recursion on $i < \operatorname{otp}(C_{\delta})$, as follows:

• if $\delta_{i+1} < \delta_i + \theta$, then set $x_i^{\delta} := \{\delta_i\}$.

• if $\delta_{i+1} \geq \delta_i + \theta$, then set $x_i^{i} := y_{\gamma,\epsilon,\tau}$, for $\gamma := \delta_i$, $\epsilon := \operatorname{otp}(\bigcup_{i' < i} x_{i'}^{\delta})$, and $\tau := h_{\delta}(\delta_i)$. In particular, $h(\operatorname{otp}(\bigcup_{i' \leq i} x_{i'}^{\delta}) - 1) = h_{\delta}(\delta_i)$.

Finally, let $C_{\delta}^{\bullet} := \bigcup_{i < \operatorname{otp}(C_{\delta})} x_i^{\delta}$, so that C_{δ}^{\bullet} is a club in δ . Note that $\operatorname{otp}(C_{\delta}) \leq \xi$, since $\operatorname{otp}(C_{\delta}) \leq \xi$ and $\alpha + \beta < \xi$ for all $(\alpha, \beta) \in \theta \times \xi$. Thus, to see that $\vec{C}^{\bullet} := \langle C_{\delta}^{\bullet} \mid \delta \in S \rangle$ and h are as sought, let D be a club in κ . By possibly shrinking D, we may assume that every element of D is an indecomposable ordinal greater than θ .

Pick $\delta \in S$ such that for every $\tau < \theta$, the following set is cofinal in δ :

$$B_{\tau} := \{ \beta < \delta \mid h_{\delta}(\beta) = \tau \& \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T \}$$

Let $\tau < \theta$ and let $\beta \in B_{\tau}$. Pick $i < \operatorname{otp}(C_{\delta})$ such that $\beta = \delta_i$. Put $\beta' := \max(x_i^{\delta})$ so that $\beta \leq \beta' < \delta_{i+1} = \min(C_{\delta}^{\bullet} \setminus (\beta'+1))$. Since $\delta_{i+1} \in D$, we know that $\delta_i + \theta < \delta_{i+1}$. Consequently,

$$h(\operatorname{otp}(C^{\bullet}_{\delta} \cap \beta')) = h(\operatorname{otp}(\bigcup_{i' \le i} x^{\delta}_{i'}) - 1) = h_{\delta}(\delta_i) = \tau,$$

as sought.

Remark 4.6. The preceding lemma should not be interpreted as saying that $\Theta_1(\ldots)$ and $\Theta_2(\ldots)$ are essentially the same, since the move from \vec{C} to $\vec{C^{\bullet}}$ may lead to the loss of coherence features of \vec{C} . In addition, the above lemma is limited to $\sigma = 1$, though a simple tweak yields that if \vec{C} is a witness for $\operatorname{CG}_{\xi}(S, T, \sigma, \langle J_{\delta} \mid \delta \in S \rangle)$ with $\sigma \leq \omega, \sigma < \theta$, and $\operatorname{nacc}(\delta) \in J_{\delta}$ for all $\delta \in S$, then a $\vec{C^{\bullet}}$ may be cooked-up to satisfy $\theta \in \Theta_1(\vec{C^{\bullet}}, T, \sigma)$.

4.1. Using colourings. We now introduce two colouring principles from [IR22a] which we shall use in this subsection. As explained in [IR22a, Remark 8.2], these principles are a spin-off of Sierpiński's *onto mapping principle*.

Definition 4.7 ([IR22a]). Let J be an ideal over λ , and $\theta \leq \lambda$ be some cardinal.

• onto (J,θ) asserts the existence of a colouring $c : [\lambda]^2 \to \theta$. such that for every $B \in J^+$, there is an $\eta < \lambda$ such that

$$c[\{\eta\} \circledast B] = \theta;$$

• unbounded (J, θ) asserts the existence of an upper-regressive colouring $c : [\lambda]^2 \to \theta$ such that for every $B \in J^+$, there is an $\eta < \lambda$ such that

$$\operatorname{otp}(c[\{\eta\} \circledast B]) = \theta.$$

Our first application of which will make use of the following pumping up result.

Fact 4.8 ([IR22b, §4]). Let $\theta \leq \lambda$ be a pair of infinite cardinals, with λ regular.

(1) If $onto(J^{bd}[\lambda], \theta)$ holds, then there exists a colouring $c : [\lambda]^2 \to \theta$ such that for every λ -complete ideal J on some ordinal δ of cofinality λ and every map $\psi : \delta \to \lambda$ satisfying $sup(\psi[B]) = \lambda$ for all $B \in J^+$, the following holds. For all $B \in J^+$, there exists an $\eta < \lambda$ such that

 $\{\tau < \theta \mid \{\beta \in B \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau\} \in J^+\} = \theta.$

(2) If unbounded($J^{bd}[\lambda], \theta$) holds, then there exists a colouring $c : [\lambda]^2 \to \theta$ such that for every λ -complete ideal J on some ordinal δ of cofinality λ and every map $\psi : \delta \to \lambda$ satisfying $\sup(\psi[B]) = \lambda$ for all $B \in J^+$, the following holds. For all $B \in J^+$, there exists an $\eta < \lambda$ such that

$$otp(\{\tau < \theta \mid \{\beta \in B \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau\} \in J^+\}) = \theta.$$

Theorem 4.9. Suppose that \vec{C} witnesses $\operatorname{CG}_{\xi}(S,T,\sigma,\vec{J})$ with $S \subseteq E_{\lambda}^{\kappa}$, and that $\theta \leq \lambda$ is infinite.

- (1) If onto $(J^{\mathrm{bd}}[\lambda], \theta)$ holds and $\xi = \lambda$, then $\theta \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$;
- (2) If onto($J^{\mathrm{bd}}[\lambda], \theta$) holds, then $\theta \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$;
- (3) If unbounded($J^{bd}[\lambda], \theta$) holds and $\theta < \lambda$, then $\theta \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$.

Proof. For every $\delta \in S$, fix a club e_{δ} in δ of ordertype λ . In case that $\xi = \lambda$, moreover set $e_{\delta} := C_{\delta}$. As J_{δ} is a λ -complete ideal on δ extending $J^{\mathrm{bd}}[\delta]$, once we define $\psi_{\delta} : \delta \to \lambda$ via $\psi_{\delta}(\beta) := \mathrm{otp}(e_{\delta} \cap \beta)$, then $\mathrm{sup}(\psi_{\delta}[B]) = \lambda$ for every $B \in (J_{\delta})^+$.

(1) and (2): Suppose that $onto(J^{bd}[\lambda], \theta)$ holds, and fix a colouring $c : [\lambda]^2 \to \theta$ as in Fact 4.8(1).

Claim 4.9.1. There exists an $\eta < \lambda$ such that, for every club $D \subseteq \kappa$, there exists $a \ \delta \in S$, such that, for every $\tau < \theta$:

$$\{\beta < \delta \mid \eta < \psi_{\delta}(\beta) \& c(\eta, \psi_{\delta}(\beta)) = \tau \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} \in J_{\delta}^{+}.$$

Proof. Suppose not. For every $\eta < \lambda$, fix a counterexample club $D_{\eta} \subseteq \kappa$. Let $D := \bigcap_{\eta < \lambda} D_{\eta}$. By the choice of \vec{C} , let us now pick $\delta \in S$ such that the following set is in J_{δ}^+ :

$$B := \{ \beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \}.$$

Recalling that c was given by Fact 4.8(1), there is an $\eta < \lambda$ such that

$$\{\tau < \theta \mid \{\beta \in B \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau\} \in (J_{\delta})^+\} = \theta.$$

 \dashv

However, as $D \subseteq D_{\eta}$, this contradicts the choice of D_{η} .

Let $\eta < \lambda$ be given by the preceding claim. Choose $\vec{h} = \langle h_{\delta} : \delta \to \theta \mid \delta \in S \rangle$ satisfying $h_{\delta}(\beta) = c(\eta, \psi_{\delta}(\beta))$ for every $\delta \in S$ and $\beta < \delta$ such that $\eta < \psi_{\delta}(\beta)$). Then \vec{h} witnesses that $\theta \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$. Furthermore, in the special case that \vec{C} is λ -bounded, any map $h : \kappa \to \theta$ satisfying $h(\bar{\beta}) = c(\eta, \bar{\beta})$ for every $\bar{\beta} \in (\eta, \lambda)$ witnesses that $\theta \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$.

(3) Suppose that unbounded($J^{\mathrm{bd}}[\lambda], \theta$) holds with $\theta < \lambda$, and fix a colouring $c : [\lambda]^2 \to \theta$ as in Fact 4.8(2). For every club $D \subseteq \kappa$, for all $\delta \in S$ and $\eta < \lambda$, denote $D(\eta, \delta) := \{\tau < \theta \mid \{\beta < \delta \mid \eta < \psi_{\delta}(\beta) \& c(\eta, \psi_{\delta}(\beta)) = \tau \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} \in J_{\delta}^+\}.$

Claim 4.9.2. There exists an $\eta < \lambda$ such that, for every club $D \subseteq \kappa$, there exists $a \ \delta \in S$, such that $|D(\eta, \delta)| = \theta$.

Proof. Suppose not. For every $\eta < \lambda$, fix a counterexample club $D_{\eta} \subseteq \kappa$. Let $D := \bigcap_{\eta < \lambda} D_{\eta}$. By the choice of \vec{C} , let us now pick $\delta \in S$ such that the following set is in J_{δ}^+ :

$$B := \{ \beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \}.$$

Recalling that c was given by Fact 4.8(2), there is an $\eta < \lambda$ such that

 $\operatorname{otp}(\{\tau < \theta \mid \{\beta \in B \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau\} \in (J_{\delta})^+\}) = \theta.$

So $|D(\eta, \delta)| = \theta$. However, as $D \subseteq D_{\eta}$, and D_{η} was chosen so that $|D_{\eta}(\eta, \delta)| < \theta$, we reach a contradiction.

We fix from hereon an $\eta < \lambda$ as given by the previous claim, and for simplicity of notation, for $D \subseteq \kappa$ a club and $\delta \in S$, we denote $D(\delta) := D(\eta, \delta)$.

Claim 4.9.3. There exists a club $D^* \subseteq \kappa$ such that for every club $D \subseteq D^*$, there exists $\delta \in S$ such that $D(\delta) = D^*(\delta)$ and this set has size θ .

Proof. Suppose this is not so. In that case, we can construct a \subseteq -decreasing sequence $\langle D_i | i \leq \theta^+ \rangle$ of clubs in κ as follows:

- (i) $D_0 := \kappa;$
- (ii) $D_{i+1} \subseteq D_i$ is some club such that for every $\delta \in S$, either $|D(\delta)| < \theta$ or $D_{i+1}(\delta) \subsetneq D_i(\delta)$;
- (iii) for $i \in \operatorname{acc}(\theta^+ + 1)$, $D_i := \bigcap_{i' < i} D_{i'}$.

Since D_{θ^+} is again a club in κ , we may fix a $\delta \in S$ such that $|D_{\theta^+}(\delta)| = \theta$. In particular, for every $i \leq \theta^+$, $|D_i(\delta)| = \theta$, and hence, by the construction, $\langle D_i(\delta) | i \leq \theta^+ \rangle$ must be a strictly \subseteq -decreasing sequence of subsets of $D_0(\delta)$, contradicting the fact that $|D_0(\delta)| = \theta$.

Let $D^* \subseteq \kappa$ be given by the preceding claim. Then any sequence $\vec{h} = \langle h_{\delta} : \delta \to \theta \mid \delta \in S \rangle$ satisfying that for all $\delta \in S$ and $\beta < \delta$ with $\psi_{\delta}(\beta) > \eta$,

$$h_{\delta}(\beta) = \operatorname{otp}(c(\eta, \psi_{\delta}(\beta)) \cap D^*(\delta))$$

witnesses that $\theta \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$.

We now move on to the case of normal ideals. We first need an analogue of Fact 4.8. In what follows, for a set of ordinals A, its *collapsing map* is the unique function $\psi : A \to \operatorname{otp}(A)$ satisfying $\psi(\alpha) = \operatorname{otp}(A \cap \alpha)$ for all $\alpha \in A$.

Theorem 4.10. Suppose that λ is a regular uncountable cardinal, and $\theta \leq \lambda$ is a cardinal.

(1) If unbounded(NS_{λ}, θ) holds, then there exists a colouring $c : [\lambda]^2 \to \theta$ such that for every λ -complete normal ideal J on some ordinal δ of cofinality λ , for every club A in δ of ordertype λ , for its collapsing map $\psi : A \to \lambda$ the following holds. For all $B \in J^+$, there exists an $\eta < \lambda$ such that

 $otp(\{\tau < \theta \mid \{\beta \in B \cap A \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau\} \in J^+\}) = \theta.$

(2) If onto(NS_{λ}, θ) holds, then there exists a colouring $c : [\lambda]^2 \to \theta$ then for every normal λ -complete ideal J on some ordinal δ of cofinality λ , for every

club A in δ of ordertype λ , for its collapsing map $\psi : A \to \lambda$ the following holds. For all $B \in J^+$, there exists an $\eta < \lambda$ such that

$$\{\tau < \theta \mid \{\beta \in B \cap A \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau\} \in J^+\} = \theta.$$

Proof. Clauses (1) and (2) follow from [IR22a, Proposition 2.25]. Since the proof of Clause (2) was omitted in [IR22a], we give it here.

(2) Given a normal λ -complete ideal J on some ordinal δ of cofinality λ , and a club A in δ of ordertype λ , it is the case that $A \in J^*$, since J is normal. Let ψ denote the collapsing map of A. For all $B \subseteq \delta$, $\eta < \lambda$ and $\tau < \theta$, denote

$$B^{\eta,\tau} := \{ \beta \in B \cap A \mid \eta < \psi(\beta) \& c(\eta, \psi(\beta)) = \tau \}.$$

Suppose onto(NS_{λ}, θ) holds, and fix a witnessing colouring $c : [\lambda]^2 \to \theta$. Towards a contradiction, suppose that there exists $B \in J^+$ such that, for every $\eta < \lambda$, there is a $\tau_{\eta} < \theta$ such that $B^{\eta,\tau_{\eta}} \in J$. As J is normal, $E := \psi^{-1}[\Delta_{\eta < \lambda}(\lambda \setminus \psi[B^{\eta,\tau_{\eta}}]]$ is in J^* . Note that

$$E = \{ \beta \in A \mid \forall \eta < \psi(\beta) \ (\beta \notin B^{\eta, \tau_{\eta}}) \}.$$

As $E \in J^*$ and $B \in J^+$, $B \cap E \in J^+$, so since J is normal, $\psi[B \cap E]$ is stationary. It thus follows from the choice of c that we may pick $\eta < \lambda$ such that $c[\{\eta\} \circledast \psi[B \cap E]] = \theta$. Find $\beta \in B \cap E$ such that $\psi(\beta) > \eta$ and $c(\eta, \psi(\beta)) = \tau_{\eta}$. Then $\beta \in B^{\eta, \tau_{\eta}}$, contradicting the fact that $\beta \in E$.

Theorem 4.11. Suppose that λ is a regular uncountable cardinal, \vec{C} witnesses $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$ with $S \subseteq E_{\lambda}^{\kappa}$ and for every $\delta \in S$, J_{δ} is a normal λ -complete ideal on δ extending $J^{\operatorname{bd}}[\delta]$. Then:

- (1) If onto(NS_{λ}, θ) holds and $\xi = \lambda$, then $\theta \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$;
- (2) If onto(NS_{λ}, θ) holds, then $\theta \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$;
- (3) If unbounded(NS_{λ}, θ) holds and $\theta < \lambda$, then $\theta_2 \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$.

Proof. Write \vec{C} as $\langle C_{\delta} | \delta \in S \rangle$. For each $\delta \in S$, if $otp(C_{\delta}) = \lambda$, then set $A_{\delta} := C_{\delta}$. Otherwise, just let A_{δ} be some club in δ of ordertype λ . Then, let $\psi_{\delta} : A_{\delta} \to \lambda$ be the corresponding collapsing map. We can now repeat the proof of Theorem 4.9 except that we use Theorem 4.10 instead of Fact 4.8.

4.2. Maintaining coherence. By Theorem 4.9, onto $(J^{\mathrm{bd}}[\lambda], \theta)$ implies that $\theta \in \Theta_1(\vec{C}, T)$. In contrast, unbounded $(J^{\mathrm{bd}}[\lambda], \theta)$ gives $\theta \in \Theta_2(\vec{C}, T)$, and then Lemma 4.5 only yields another *C*-sequence \vec{C}^{\bullet} such that $\theta \in \Theta_1(\vec{C}^{\bullet}, T)$.

In the next theorem, we combine the two results carefully in order to obtain a C-sequence \vec{C}^{\bullet} with $\theta \in \Theta_1(\vec{C}^{\bullet}, T)$ while maintaining some coherence features of the original sequence \vec{C} .

Theorem 4.12. Suppose that $\theta < \lambda < \kappa$ are infinite cardinals, unbounded $(J^{bd}[\lambda], \theta)$ holds, and S is a stationary subset of E_{λ}^{κ} .

For every C-sequence $\vec{C} = \langle C_{\delta} \mid \delta < \kappa \rangle$ such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}_{\lambda}(S,T)$, there exists a corresponding C-sequence $\vec{C}^{\bullet} = \langle C_{\delta}^{\bullet} \mid \delta < \kappa \rangle$ such that:

- $\vec{C}^{\bullet} \upharpoonright S$ is λ -bounded;
- If \vec{C} is weakly coherent, then so is \vec{C}^{\bullet} ;
- For every infinite cardinal $\chi \in [\theta, \kappa)$, if \vec{C} is $\chi \sqsubseteq$ -coherent, then so is \vec{C}^{\bullet} ;
- $\theta \in \Theta_1(\vec{C}^{\bullet} \upharpoonright S, T).$

Proof. Without loss of generality, $0 \in C_{\delta}$ for all nonzero $\delta < \kappa$. For every $\delta < \kappa$, denote $\xi_{\delta} := \operatorname{otp}(C_{\delta})$, let $\psi_{\delta} : C_{\delta} \to \xi_{\delta}$ be the collapsing map of C_{δ} , and let $\langle \delta_i |$ $i < \xi_{\delta}$ denote the increasing enumeration of C_{δ} so that $\psi_{\delta}(\delta_i) = i$ for every $i < \xi_{\delta}$.

Fix a colouring $c: [\lambda]^2 \to \theta$ as in Fact 4.8(2). For every club $D \subseteq \kappa$, for all $\delta \in S$ and $\eta < \lambda$, denote

$$D(\eta, \delta) := \{ \tau < \theta \mid \sup\{\beta < \delta \mid c(\eta, \psi_{\delta}(\beta)) = \tau \& \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T \} = \delta \}.$$

By Claims 4.9.2 and 4.9.3, we may pick $\eta^* < \lambda$ and a club $D^* \subseteq \kappa$ such that for every club $D \subseteq D^*$, there exists $\delta \in S$ such that $D(\eta^*, \delta) = D^*(\eta^*, \delta)$ and this set has size θ . By possibly shrinking D^* , we may assume that D^* consists of indecomposable ordinals, and that $\min(D^*) > \theta$.

For every $\delta \in S$, since $cf(\delta) = \lambda > \theta$, the set

$$N_{\delta} := \{\beta < \delta \mid c(\eta^*, \psi_{\delta}(\beta)) \notin D^*(\eta^*, \delta) \& \min(C_{\delta} \setminus (\beta + 1)) \in D^* \cap T\}$$

is bounded in δ . So, by one more stabilization argument, we may fix an $\varepsilon < \lambda$ such that every club $D \subseteq D^*$, there exists $\delta \in S$ such that $D(\eta^*, \delta) = D^*(\eta^*, \delta)$, $|D^*(\eta^*, \delta)| = \theta$, and also

$$\sup(\{\psi_{\delta}(\beta) \mid \beta \in N_{\delta}\}) = \varepsilon.$$

For all $\delta \in \operatorname{acc}(\kappa)$ and $i \leq \xi_{\delta}$, denote:

$$T^{i}_{\delta} := \{ c(\eta^{*}, j) \mid \varepsilon < j < i, \ \eta^{*} < j, \ \delta_{j+1} \in D^{*} \cap T \}.$$

Fix a surjection $h: \kappa \to \theta$ and a sequence of sets $\langle y_{\gamma,\epsilon,\tau} \mid (\gamma,\epsilon,\tau) \in \kappa \times \kappa \times \theta \rangle$ as in the proof of Lemma 4.5. We now construct the new C-sequence $\vec{C} \bullet = \langle C_{\delta} \bullet \rangle$ $\delta < \kappa \rangle$, as follows. Set $C_0^{\bullet} := \emptyset$ and $C_{\gamma+1}^{\bullet} := \{\gamma\}$ for every $\gamma < \kappa$. Next, given $\delta \in \operatorname{acc}(\kappa)$, construct a sequence $\langle x_{\delta}^i \mid i < \xi_{\delta} \rangle$ by recursion on $i < \xi_{\delta}$, as follows:

• If $\delta_{i+1} \notin D^*$, then set $x_{\delta}^i := \{\delta_i\}$.

▶ If $\delta_{i+1} \in D^*$, then in particular, $\delta_{i+1} \geq \delta_i + \theta$, so we set $x^i_{\delta} := y_{\gamma,\epsilon,\tau}$, for $\gamma := \delta_i, \ \epsilon := \operatorname{otp}(\bigcup_{i' < i} x_{\delta}^{i'}), \ \text{and}$

$$\tau := \operatorname{otp}(c(\eta^*, i) \cap T^i_{\delta}).$$

Note that, for every $\delta \in \operatorname{acc}(\kappa)$, $C_{\delta} \subseteq C^{\bullet}_{\delta}$, and also $\operatorname{acc}(C^{\bullet}_{\delta}) \cap E^{\kappa}_{\geq \theta} = \operatorname{acc}(C_{\delta})$, since $y_{\gamma,\epsilon,\tau} \subseteq E_{<\theta}^{\kappa}$ for every $(\gamma,\epsilon,\tau) \in \kappa \times \kappa \times \theta$. In addition, for every $\delta \in \operatorname{acc}(\kappa)$, if $\alpha + \beta < \xi_{\delta}$ for all $(\alpha, \beta) \in \theta \times \xi_{\delta}$, then $\operatorname{otp}(C^{\bullet}_{\delta}) = \xi_{\delta}$. In particular, $\operatorname{otp}(C^{\bullet}_{\delta}) = \lambda$ for all $\delta \in S$.

Claim 4.12.1. Let $\chi \in [\theta, \kappa)$ be an infinite cardinal. If \vec{C} is ${}_{\chi}\sqsubseteq$ -coherent, then so is \vec{C}^{\bullet} .

Proof. Suppose that \vec{C} is $\chi \sqsubseteq$ -coherent. Let $\delta < \kappa$ and $\bar{\delta} \in \operatorname{acc}(C^{\bullet}_{\delta}) \cap E^{\kappa}_{\geq \chi}$; we need to verify that $C^{\bullet}_{\delta} \cap \overline{\delta} \sqsubseteq C^{\bullet}_{\delta}$. As $\operatorname{acc}(C^{\bullet}_{\delta}) \cap E^{\kappa}_{\geq \theta} = \operatorname{acc}(C_{\delta})$ and $\chi \geq \theta$, we infer that $\overline{\delta} \in \operatorname{acc}(C_{\delta})$, so by ${}_{\chi} \sqsubseteq$ -coherence of \overline{C} , $C_{\delta} \cap \overline{\delta} \sqsubseteq C_{\delta}$. It follows that:

- $\langle \bar{\delta}_i \mid i < \xi_{\bar{\delta}} \rangle = \langle \delta_i \mid i < \xi_{\bar{\delta}} \rangle$, $\langle T^i_{\bar{\delta}} \mid i < \xi_{\bar{\delta}} \rangle = \langle T^i_{\delta} \mid i < \xi_{\bar{\delta}} \rangle$, and hence $\langle x^i_{\bar{\delta}} \mid i < \xi_{\bar{\delta}} \rangle = \langle x^i_{\delta} \mid i < \xi_{\bar{\delta}} \rangle$,

so $C^{\bullet}_{\delta} \cap \bar{\delta} = \bigcup_{i < f_{\pi}} x^i_{\delta} = \bigcup_{i < f_{\pi}} x^i_{\bar{\delta}} = C^{\bullet}_{\bar{\delta}}$, as sought.

 \dashv

Claim 4.12.2. If \vec{C} is weakly coherent, then so is \vec{C}^{\bullet} .

Proof. Towards a contradiction, suppose that \vec{C} is weakly coherent, but $\vec{C^{\bullet}}$ is not. Fix the least $\alpha < \kappa$ such that $|\{C^{\bullet}_{\delta} \cap \alpha \mid \delta < \kappa\}| = \kappa$. So we may fix a cofinal subset Δ of $\operatorname{acc}(\kappa)$ such that:

- (1) $\delta \mapsto C^{\bullet}_{\delta} \cap \alpha$ is injective over Δ , but
- (2) $\delta \mapsto C_{\delta} \cap \alpha$ is constant over Δ .

Fix $\gamma < \alpha$ such that $\sup(C_{\delta} \cap \alpha) = \gamma$ for all $\delta \in \Delta$. By minimality of α , and by possibly shrinking Δ further, we may also assume that

(3) $\delta \mapsto C^{\bullet}_{\delta} \cap \gamma$ is constant over Δ .

It thus follows that the map $\delta \mapsto C^{\bullet}_{\delta} \cap [\gamma, \alpha)$ is injective over Δ . However, for every $\delta \in \Delta$, $C^{\bullet}_{\delta} \cap [\gamma, \alpha)$ is equal to $y_{\gamma, \epsilon, \tau} \cap \alpha$, for $\epsilon := \operatorname{otp}(C^{\bullet}_{\delta} \cap \gamma)$ and some $\tau < \theta$. Recalling Clause (3), there exists an $\epsilon < \kappa$ such that:

$$\{C^{\bullet}_{\delta} \cap [\gamma, \alpha) \mid \delta \in \Delta\} \subseteq \{y_{\gamma, \epsilon, \tau} \cap \alpha \mid \tau < \theta\},\$$

contradicting the fact that the set on the right hand size has size $\leq \theta < \kappa$. \dashv

Finally, to see that $\theta \in \Theta_1(\vec{C} \cap S, T)$, let D be a club in κ . By possibly shrinking D, we may assume that $D \subseteq D^*$. Pick $\delta \in S$ such that $D(\eta^*, \delta) = D^*(\eta^*, \delta)$, $|D^*(\eta^*, \delta)| = \theta$, and also

$$\sup(\{\psi_{\delta}(\beta) \mid \beta \in N_{\delta}\}) = \varepsilon.$$

For any $D' \in \{D, D^*\},\$

 $D'(\eta^*, \delta) = \{ \tau < \theta \mid \sup\{i < \lambda \mid c(\eta^*, i) = \tau \& \delta_{i+1} \in D' \cap T \} = \lambda \}.$

So, since $D(\eta^*, \delta) = D^*(\eta^*, \delta)$, the definition of ε implies that

$$D(\eta^*, \delta) = \{ c(\eta^*, j) \mid \varepsilon < j < \lambda, \ \eta^* < j, \ \delta_{j+1} \in D^* \cap T \} = T_{\delta}^{\lambda}.$$

In particular, $|T_{\delta}^{\lambda}| = \theta$. Now, given a prescribed colour τ^* and some $\alpha < \delta$, we shall find a $\beta^* \in C_{\delta}^{\bullet}$ above α such that $\min(C_{\delta}^{\bullet} \setminus (\beta^* + 1)) \in D \cap T$ and $h(C_{\delta}^{\bullet} \cap \beta^*) = \tau^*$. Fix the unique $\tau \in T_{\delta}^{\lambda}$ such that $\operatorname{otp}(T_{\delta}^{\lambda} \cap \tau) = \tau^*$. Note that for a tail of $i < \lambda$, $T_{\delta}^{\lambda} \cap \tau = T_{\delta}^{i} \cap \tau$.

As $\tau \in D(\eta^*, \delta)$, there are cofinally many $\beta \in C_{\delta}$ such that $\min(C_{\delta} \setminus (\beta + 1)) \in D \cap T$, $\eta^* < \psi_{\delta}(\beta)$ and $c(\eta^*, \psi_{\delta}(\beta)) = \tau$. So, we may find a large enough $i < \lambda$ such that:

- $\delta_{i+1} \in D \cap T$,
- $\eta^* < i$,
- $c(\eta^*, i) = \tau$,
- $\alpha < \delta_i$, and
- $T^{\lambda}_{\delta} \cap \tau = T^{i}_{\delta} \cap \tau.$

So $\operatorname{otp}(c(\eta^*, i) \cap T^i_{\delta}) = \operatorname{otp}(\tau \cap T^i_{\delta}) = \operatorname{otp}(T^{\lambda}_{\delta} \cap \tau) = \tau^*$. Put $\beta^* := \max(x^i_{\delta})$ so that $\delta_i \leq \beta^* < \delta_{i+1} = \min(C^{\bullet}_{\delta} \setminus (\beta^* + 1))$. Since $\delta_{i+1} \in D \subseteq D^*$, we know that $x^i_{\delta} = y_{\gamma,\epsilon,\tau^*}$, for $\gamma := \delta_i$ and $\epsilon := \operatorname{otp}(\bigcup_{i' < i} x^{i'}_{\delta})$. Consequently,

$$h(\operatorname{otp}(C^{\bullet}_{\delta} \cap \beta^*)) = h(\operatorname{otp}(\bigcup_{i' \le i} x^{\delta}_{i'}) - 1) = \tau^*,$$

as sought.

By [IR22a, Proposition 2.18 and Lemma 8.4], in Gödel's constructible universe L, for every weakly compact cardinal λ that is not ineffable, unbounded $(J^{\text{bd}}[\lambda], \theta)$ fails for every cardinal $\theta \in [3, \lambda]$, but $onto(NS_{\lambda}, \lambda)$ holds. Thus, it is easier to get unbounded (J, θ) with $J := NS_{\lambda}$ than with $J := J^{\text{bd}}[\lambda]$. In the upcoming

theorem, the hypothesis of unbounded $(J^{\mathrm{bd}}[\lambda], \theta)$ from Theorem 4.12 is reduced to unbounded $(\mathrm{NS}_{\lambda}, \theta)$ at the cost of requiring $\vec{C} \upharpoonright S$ to witness $\mathrm{CG}_{\lambda}(S, T, 1, \langle \mathrm{NS}_{\delta} \mid \delta \in S \rangle)$.

Theorem 4.13. Suppose that $\theta < \lambda < \kappa$ are infinite cardinals, unbounded(NS_{λ}, θ) holds, and S is a stationary subset of E_{λ}^{κ} .

For every C-sequence $\vec{C} = \langle C_{\delta} \mid \delta < \kappa \rangle$ such that $\vec{C} \upharpoonright S$ witnesses $\operatorname{CG}_{\lambda}(S, T, 1, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$, there exists a corresponding C-sequence $\vec{C}^{\bullet} = \langle C^{\bullet}_{\delta} \mid \delta < \kappa \rangle$ such that:

- $\vec{C}^{\bullet} \upharpoonright S$ is λ -bounded;
- If \vec{C} is weakly coherent, then so is \vec{C}^{\bullet} ;
- For every infinite cardinal $\chi \in [\theta, \kappa)$, if \vec{C} is $\chi \sqsubseteq$ -coherent, then so is \vec{C}^{\bullet} ;
- $\theta \in \Theta_1(\vec{C^{\bullet}} \upharpoonright S, T).$

Proof. For every $\delta < \kappa$, let $\psi_{\delta} : C_{\delta} \to \operatorname{otp}(C_{\delta})$ denote the collapsing map of C_{δ} . Let $c : [\kappa]^2 \to \theta$ be a colouring given by Theorem 4.10(1).

Claim 4.13.1. There exists an $\eta < \lambda$ such that, for every club $D \subseteq \kappa$, there exists $a \ \delta \in S$, such that the following set has size θ :

$$D(\eta, \delta) := \{ \tau < \theta \mid \sup\{\beta < \delta \mid c(\eta, \psi_{\delta}(\beta)) = \tau \& \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T \} = \delta \}.$$

Proof. Suppose not. For every $\eta < \lambda$, fix a counterexample club $D_{\eta} \subseteq \kappa$. Let $D := \bigcap_{\eta < \lambda} D_{\eta}$. By the hypothesis on $\vec{C} \upharpoonright S$, let us now pick $\delta \in S$ such that the following set is stationary in δ :

$$B := \{\beta < \delta \mid \min(C_{\delta} \setminus (\beta + 1)) \in D \cap T\}.$$

By the choice of c, there is an $\eta < \lambda$ such that

$$otp(\{\tau < \theta \mid \{\beta \in B \mid \eta < \psi_{\delta}(\beta) \& c(\eta, \psi_{\delta}(\beta)) = \tau\} \in (NS_{\delta})^+\}) = \theta.$$

In particular, $|D(\eta, \delta)| = \theta$, contradicting the fact that $D \subseteq D_{\eta}$.

The rest of the proof is now identical to that of Theorem 4.12.

Remark 4.14. Since $\operatorname{nacc}(\delta) \in \operatorname{NS}_{\delta}$ for all $\delta \in S$, for every $\sigma \leq \omega$ such that $\sigma < \theta$, if $\vec{C} \upharpoonright S$ moreover witnesses $\operatorname{CG}_{\lambda}(S, T, \sigma, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$, then the preceding proof may be slightly tweaked to secure that $\theta \in \Theta_1(\vec{C} \bullet \upharpoonright S, T, \sigma)$. The first change is to require that the surjection $h : \kappa \to \theta$ satisfies that for every $\epsilon < \kappa$, for every $\tau < \theta$, there exists $\iota \in (\epsilon, \epsilon + \theta)$ such that $\{h(\iota + \varsigma) \mid \varsigma \leq \sigma\} = \{\tau\}$. The second change is to impose $x^i_{\delta} = \{\delta_i\}$ for all $i \in \operatorname{nacc}(\xi_{\delta})$. The details are left to the reader.

4.3. **Applications.** We now utilize the results from [IR22a, IR22b] in order to partition club-guessing sequences.

Fact 4.15 ([IR22a]). Suppose that λ is regular and uncountable.

Any of the following implies that $onto(J^{bd}[\lambda], \theta)$ holds:

- (1) $\theta = \lambda = \aleph_1 = \operatorname{non}(\mathcal{M});$
- (2) $\theta = \lambda$ is a successor cardinal, and (λ) holds;
- (3) $\theta = \lambda$ and $\Diamond(T)$ holds for a stationary $T \subseteq \lambda$ that does not reflect at regulars;
- (4) $\theta < \lambda$ and $\lambda \not\rightarrow [\lambda]^2_{\theta}$ holds;
- (5) $\theta < \lambda$ is regular and unbounded $(J^{bd}[\lambda], \lambda)$ holds.

Fact 4.16 ([IR22b]). Suppose that λ is regular and uncountable, and $\theta \leq \lambda$. Any of the following implies that unbounded($J^{bd}[\lambda], \theta$) holds:

- (1) λ admits a nontrivial C-sequence in the sense of [Tod07, Definition 6.3.1];
- (2) $\Box(\lambda, <\mu)$ holds for some $\mu < \lambda$;
- (3) λ is not greatly Mahlo;
- (4) λ is not weakly compact in L;
- (5) λ is not weakly compact, and $\theta = \omega$;
- (6) λ is not strongly inaccessible, and $\theta = \log_2(\lambda)$.

Corollary 4.17. Suppose that λ is a regular uncountable cardinal, and \vec{C} witnesses $CG_{\lambda}(S,T,\sigma,\vec{J})$ with $S \subseteq E_{\lambda}^{\kappa}$.

- (1) If $\lambda = \lambda^{<\lambda}$ is a successor cardinal, then $\lambda \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$;
- (2) If $\lambda = \theta^+$ and θ is regular, then $\theta \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$;
- (3) If λ is not Mahlo and $\Diamond(\lambda)$ holds, then $\lambda \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$;
- (4) If λ is not greatly Mahlo then $\theta \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$ for every cardinal $\theta < \lambda$;
- (5) If λ is not strongly inaccessible, then $\log_2(\lambda) \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$;
- (6) If λ is not weakly compact, then $\omega \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$.

Fact 4.18 ([IR22a]). Let λ be a regular uncountable cardinal.

- (1) if $\diamondsuit^*(\lambda)$ holds, then so does onto(NS_{λ}, λ);
- (2) If λ admits an amenable C-sequence, then onto(NS_{λ}, θ) holds for all regular $\theta < \lambda$;
- (3) unbounded(NS_{λ}, ω) holds iff λ is not ineffable.

Corollary 4.19. Suppose that λ is a regular uncountable cardinal, and \vec{C} witnesses $CG_{\lambda}(S,T,\sigma,\vec{J})$ with $S \subseteq E_{\lambda}^{\kappa}$ and \vec{J} is a sequence of normal ideals.

- (1) If $\diamond^*(\lambda)$ holds, then $\lambda \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$;
- (2) If λ admits an amenable C-sequence, then $\theta \in \Theta_1(\vec{C}, T, \sigma, \vec{J})$ for all regular $\theta < \lambda$;
- (3) If λ is not ineffable, then $\omega \in \Theta_2(\vec{C}, T, \sigma, \vec{J})$.

In [Moo08], the (weak) club-guessing principle \mathcal{V} was shown to give rise to a C-sequence \vec{C} over ω_1 for which the corresponding object $\mathcal{T}(\rho_0^{\vec{C}})$ is a special Aronszajn tree of pathological nature. In the terminology developed in this paper, the key features of \vec{C} sufficient for the construction are that \vec{C} be a transversal for \Box_{ω}^* and that $\omega \in \Theta_1(\vec{C}, \omega_1)$. Arguably, the higher analog of this would assert the existence of a transversal $\vec{C} = \langle C_{\delta} | \delta < \lambda^+ \rangle$ for \Box_{λ}^* such that $\log_2(\lambda) \in \Theta_1(\vec{C}, \lambda^+)$. By the next corollary, if λ is not strongly inaccessible (in particular, if $\lambda = \aleph_1$), then the existence of such a C-sequence is in fact no stronger than \Box_{λ}^* .

Corollary 4.20. Suppose that λ is a regular uncountable cardinal, and $\theta = \log_2(\lambda)$.

- (1) If \Box^*_{λ} holds and $\theta < \lambda$, then there exists a transversal \vec{C} for \Box^*_{λ} such that $\theta \in \Theta_1(\vec{C} \upharpoonright E^{\lambda^+}_{\lambda}, E^{\lambda^+}_{\lambda});$
- (2) If $\Diamond(\lambda)$ holds and λ is not Mahlo, then there exists a transversal \vec{C} for \Box^*_{λ} such that $\theta \in \Theta_1(\vec{C} \upharpoonright E^{\lambda^+}_{\lambda}, E^{\lambda^+}_{\lambda});$
- (3) If $\diamondsuit^*(\lambda)$ holds, then there exists a transversal \vec{C} for \Box^*_{λ} such that $\theta \in \Theta_1(\vec{C} \upharpoonright E_{\lambda}^{\lambda^+}, \lambda^+)$.

Proof. (1) Suppose that $\theta < \lambda$ and \Box_{λ}^{*} holds. Appeal to Theorem 3.20 to find a transversal \vec{C} for \Box_{λ}^{*} such that $\vec{C} \upharpoonright E_{\lambda}^{\lambda^{+}}$ witnesses $\operatorname{CG}_{\lambda}(E_{\lambda}^{\lambda^{+}}, E_{\lambda}^{\lambda^{+}})$. By Corollary 4.17(5), $\theta \in \Theta_{1}(\vec{C} \upharpoonright E_{\lambda}^{\lambda^{+}}, E_{\lambda}^{\lambda^{+}})$.

(2) Appeal to Fact 3.25 to pick a λ -bounded *C*-sequence $\vec{C} = \langle C_{\delta} | \delta < \lambda^+ \rangle$ such that $C \upharpoonright E_{\lambda}^{\lambda^+}$ witnesses $CG_{\lambda}(E_{\lambda}^{\lambda^+}, E_{\lambda}^{\lambda^+})$. Suppose that $\Diamond(\lambda)$ holds. In particular, $\lambda^{<\lambda} = \lambda$, so \vec{C} is trivially weakly coherent. In addition, since $\Diamond(\lambda)$ holds, Corollary 4.17(3) implies that $\lambda \in \Theta_1(E_{\lambda}^{\lambda^+}, E_{\lambda}^{\lambda^+})$.

(3) By Fact 3.25 together with Corollary 6.10 below, we may fix a λ -bounded C-sequence $\vec{C} = \langle C_{\delta} \mid \delta < \lambda^+ \rangle$ such that $\vec{C} \upharpoonright E_{\lambda}^{\lambda^+}$ witnesses $\operatorname{CG}_{\lambda}(E_{\lambda}^{\lambda^+}, \lambda^+, 1, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$. Suppose that $\Diamond^*(\lambda)$ holds. In particular, \vec{C} is trivially weakly coherent. In addition, since $\Diamond^*(\lambda)$ holds, Corollary 4.19(1) implies that $\lambda \in \Theta_1(\vec{C} \upharpoonright E_{\lambda}^{\lambda^+}, \lambda^+)$. \Box

We also record the following variation.

Corollary 4.21. Suppose that λ is a regular uncountable cardinal that is not strongly inaccessible. If \Box_{λ} holds, then it may be witnessed by a *C*-sequence \vec{C} such that $\log_2(\lambda) \in \Theta_1(\vec{C} \upharpoonright E_{\lambda}^{\lambda^+}, \lambda^+)$.

Proof. By Fact 3.4, using $(\xi, \kappa, \mu, S) := (\lambda, \lambda^+, 2, E_{\lambda}^{\lambda^+})$, and Corollary 4.17(5). \Box

4.4. Another form of partitioning. By [BR21, Theorem 3.7], if $\clubsuit(S)$ holds, then there exists a decomposition $S = \biguplus_{i < \kappa} S_i$ such that $\clubsuit(S_i)$ holds for every $i < \kappa$. We close this section by showing that this form of partitioning also holds for CG. When taken together with Theorem 2.15, this yields Solovay's decomposition theorem for cardinals greater than \aleph_1 (at the level of \aleph_1 , Solovay's theorem follows using an Ulam matrix).

Proposition 4.22. Suppose that \vec{C} witnesses $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$. Then there exists a decomposition $S = \biguplus_{i < \kappa} S_i$ such that $\vec{C} \upharpoonright S_i$ witnesses $\operatorname{CG}_{\xi}(S_i, T, \sigma, \vec{J})$ for all $i < \kappa$.

Proof. For all $\beta < i < \kappa$, let $S_i^{\beta} := \{\delta \in S \cap \operatorname{acc}(\kappa \setminus \beta) \mid \min(C_{\delta} \setminus (\beta + 1)) = i\}$. It suffices to prove that there exists a $\beta < \kappa$ for which the following set has size κ :

 $I_{\beta} := \{ i \in (\beta, \kappa) \mid \vec{C} \upharpoonright S_{i}^{\beta} \text{ witnesses } \operatorname{CG}_{\xi}(S_{i}^{\beta}, T, \sigma, \vec{J}) \}.$

So, suppose that this is not the case, and fix a sparse enough club $E \subseteq \kappa$ such that, for every $\epsilon \in E$, for every $\beta < \epsilon$, $\sup(I_{\beta}) < \epsilon$. In addition, fix a triangular matrix $\langle D_i^{\beta} | \beta < i < \kappa \rangle$ of clubs in κ such that, for all $\beta < i < \kappa$, if $i \notin I_{\beta}$, then for every $\delta \in S_i^{\beta}$,

$$\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D_i^{\beta} \cap T\} \in J_{\delta}.$$

Consider the club $D := \{\delta \in \operatorname{acc}(E) \mid \forall i < \delta \forall \beta < i(\delta \in D_i^\beta)\}$. By the choice of \vec{C} , pick $\delta \in S$ such that the following set is in J_{δ}^+ :

$$B := \{ \beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \}.$$

Claim 4.22.1. For every $\beta < \delta$, $\min(C_{\delta} \setminus (\beta + 1)) \in I_{\beta}$.

Proof. Given $\beta < \delta$, if we let $i := \min(C_{\delta} \setminus (\beta + 1))$, then $\delta \in S_i^{\beta}$, and since $D \cap \delta$ is almost included in $D_i^{\beta} \cap \delta$, it is the case that

$$\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D_i^{\beta} \cap T\} \in J_{\delta}^+,$$

so that $i \in I_{\beta}$.

Pick $\beta \in B$ and set $\epsilon := \min(E \setminus (\beta + 1))$. Since $\min(C_{\delta} \setminus (\beta + 1))$ is in $\operatorname{acc}(E)$, we infer that

$$\beta < \epsilon < \min(C_{\delta} \setminus (\beta + 1)).$$

As $\beta < \epsilon$ with $\epsilon \in E$, $\sup(I_{\beta}) < \epsilon$, contradicting the preceding claim.

5. Increasing σ

In this section we are interested in improving the quality of guessing by guessing many consecutive nonaccumulation points as in Question 1.6. As we shall see, guessing clubs relative to points of prescribed cofinality turns out be of great help for this purpose. The main result of this section reads as follows.

Corollary 5.1. Suppose $\nu < \xi \leq \kappa$ are infinite cardinals, and that $S \subseteq E_{>\nu}^{\kappa}$.

For every (possibly finite) cardinal $\sigma < \nu$, if $\operatorname{CG}_{\xi}(S, E_{>\sigma}^{\kappa} \cap E_{<\nu}^{\kappa}, 1, \vec{J})$ holds, then so does $\operatorname{CG}_{\xi}(S, \kappa, \sigma, \vec{J})$.

Proof. \blacktriangleright If σ is finite, then appeal to Theorem 5.2 below.

- ▶ If $\sigma = \omega$ and $\operatorname{CG}_{\xi}(S, E_{\sigma}^{\kappa}, 1, \vec{J})$ holds then appeal to Theorem 5.4 below.
- ▶ If $\sigma = \omega$ and $\operatorname{CG}_{\xi}(S, E_{\sigma}^{\kappa}, 1, \vec{J})$ fails, then since $S \subseteq E_{>\nu}^{\kappa}$, it follows from Proposition 2.5 that $CG_{\xi}(S, E_{\bar{\nu}}^{\kappa}, 1, \vec{J})$ holds for some cardinal $\bar{\nu}$ with $\sigma < \bar{\nu} \leq \nu$. Now appeal to Theorem 5.5(2) below with $T := \kappa$.

• If $\sigma > \omega$, then appeal to Theorem 5.5 below with $T := \kappa$.

Another key result is Theorem 5.5 below where a version of this result relative to a set T is proved.

We commence with a result that pumps up σ from 1 to any prescribed positive integer using a postprocessing^{*} function (thereby, preserving coherence features).

Theorem 5.2. Suppose $\sigma < \omega \leq \nu < \kappa$ are cardinals.

Suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, E_{<\nu}^{\kappa}, 1, \vec{J})$, with $S \subseteq E_{>\nu}^{\kappa}$. Then there exists a postprocessing^{*} function $\Phi: \mathcal{K}(\kappa) \to \overline{\mathcal{K}(\kappa)}$ such that $\langle \Phi(C_{\delta}) \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\mathcal{E}}(S, \kappa, \sigma, \vec{J})$.

Proof. Fix an auxiliary C-sequence $\vec{e} = \langle e_{\gamma} | \gamma < \kappa \rangle$ such that $otp(e_{\gamma}) = cf(\gamma)$ for every $\gamma < \kappa$. In what follows we shall use the operator Φ_D from Definition 2.12.

Claim 5.2.1. There is a club $D \subseteq \kappa$ such that for every club $E \subseteq \kappa$ there is a $\delta \in S$ such that

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{<\nu}^{\kappa} \& |\Phi_D(e_{\gamma}) \cap (E \setminus \alpha)| > \sigma\} \in J_{\delta}^+.$$

Proof. Suppose that the claim does not hold. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that for every $\delta \in S$,

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\leq \nu}^{\kappa} \& |\Phi_D(e_{\gamma}) \cap (F^D \setminus \alpha)| > \sigma\} \in J_{\delta}.$$

Let $\mu := \aleph_0$ so that $\mu \leq \nu < \kappa$. We construct now a \subseteq -decreasing sequence $\langle D_i |$ $i \leq \mu$ of clubs in κ as follows:

- (i) $D_0 := \kappa;$
- (ii) $D_{i+1} := D_i \cap F^{D_i};$ (iii) for $i \in \operatorname{acc}(\mu + 1), D_i := \bigcap_{i' < i} D_{i'}.$

Since $\mu < \kappa$, all these are clubs in κ . Finally, consider the club

$$D^* := \{ \gamma < \kappa \mid \operatorname{otp}(D_\mu \cap \gamma) = \gamma > \nu \}.$$

As
$$\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$$
 witnesses $\operatorname{CG}_{\xi}(S, E_{\leq \nu}^{\kappa}, 1, \vec{J})$, let us pick $\delta \in S$ such that
$$A := \{ \alpha < \delta \mid \min(C_{\delta} \setminus (\alpha + 1)) \in D^* \cap E_{<\nu}^{\kappa} \}$$

is in J_{δ}^+ .

For every $i < \mu$, by the choice of F^{D_i} , the following set

$$A_i := \{ \alpha < \delta \mid \gamma := \min(C_\delta \setminus (\alpha + 1)) \& \gamma \in E_{\leq \nu}^{\kappa} \& |\Phi_{D_i}(e_\gamma) \cap (F^{D_i} \setminus \alpha)| > \sigma \}$$

is in J_{δ} . As J_{δ} is $cf(\delta)$ -additive, and $cf(\delta) > \nu$, we may now fix $\alpha \in A \setminus \bigcup_{i < \mu} A_i$. Set $\gamma := \min(C_{\delta} \setminus (\alpha + 1))$, so that $\gamma \in D^* \cap E_{<\nu}^{\kappa}$. Since $\gamma \in D^* \subseteq \operatorname{acc}(D_i)$ for every $i \leq \mu$, we have that $\Phi_{D_i}(e_{\gamma}) \subseteq D_i$.

As $\gamma \in D^*$, $\operatorname{otp}(D^* \cap \gamma) = \gamma > \nu \ge \operatorname{otp}(e_{\gamma})$, so that $(D^* \cap \gamma) \setminus e_{\gamma}$ is cofinal in γ . Recursively construct $\langle (\delta^n, \beta^n) \mid n \leq \sigma \rangle$ by letting

- (i) $\delta^0 := \min(D^* \setminus (e_\gamma \cup \alpha))$, and
- (ii) $\beta^0 := \min(e_{\gamma} \setminus \delta^0)$, and
- (iii) $\delta^{n+1} := \min(D^* \setminus (e_{\gamma} \cup \beta^n))$, and (iv) $\beta^{n+1} := \min(e_{\gamma} \setminus \delta^{n+1})$.

Evidently, $\alpha \leq \delta^n < \beta^n < \delta^{n+1}$.

For every $n \leq \sigma$, denote $\beta_i^n := \sup(\beta^n \cap D_i)$, and fix a large enough $j_n < \omega$ such that $\beta_i^n = \beta_{j_n}^n$ for every integer $i \ge j_n$. Set $i^* := \max\{j_n \mid n \le \sigma\}$ which is finite as σ is finite. Altogether, for every $n < \sigma$,

$$\alpha \le \delta^n \le \beta_{i^*}^n = \beta_{i^*+1}^n \le \beta^n < \delta^{n+1}$$

It thus follows that $\{\beta_{i^*}^n \mid n \leq \sigma\}$ consists of $\sigma + 1$ many distinct elements of $\Phi_{D_{i^*}}(e_{\gamma}) \cap (D_{i^*+1} \setminus \alpha)$. But D_{i^*+1} is a subset of $F^{D_{i^*}}$ so $|\Phi_{D_{i^*}}(e_{\gamma}) \cap (F^{D_{i^*}} \setminus \alpha)| > \sigma$, and since $\gamma = \min(C_{\delta} \setminus (\alpha + 1))$, we have contradicted the fact that $\alpha \notin A_{i^*}$. +

Let $D \subseteq \kappa$ be a club as given by the preceding claim. For all $\gamma \in E_{<\nu}^{\kappa}$ and $z \in [\nu]^{<\omega}$, we define a finite subset of γ :

$$z_{\gamma} := \{ \eta \in \Phi_D(e_{\gamma}) \mid \operatorname{otp}(\Phi_D(e_{\gamma}) \cap \eta) \in z \}.$$

Now, for every $z \in [\nu]^{<\omega}$, define a postprocessing^{*} function $\Phi^z : \mathcal{K}(\kappa) \to \mathcal{K}(\kappa)$ via:

$$\Phi^{z}(x) := x \cup \{z_{\gamma} \setminus \sup(x \cap \gamma) \mid \gamma \in \operatorname{nacc}(x) \cap E_{<\nu}^{\kappa}\}$$

Claim 5.2.2. There exists $z \in [\nu]^{\sigma+1}$ such that $\vec{C^z} := \langle \Phi^z(C_\delta) \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\mathcal{E}}(S,\kappa,\sigma,\vec{J}).$

Proof. Suppose not. For each $z \in [\nu]^{\sigma+1}$ fix a counterexample E_z . Set $E := \bigcap \{E_z \mid$ $z \in [\nu]^{\sigma+1}$. Recalling the choice of D, let us now fix $\delta \in S$ for which

$$A := \{ \alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\leq \nu}^{\kappa} \& |\Phi_D(e_{\gamma}) \cap (E \setminus \alpha)| > \sigma \}$$

is in J_{δ}^+ . For every $\alpha \in A$, let $\gamma_{\alpha} := \min(C_{\delta} \setminus (\alpha + 1))$ and fix $z_{\alpha} \in [\nu]^{\sigma+1}$ such that $(z_{\alpha})_{\gamma_{\alpha}} \subseteq E \setminus \alpha$. As J_{δ} is $cf(\delta)$ -additive and $cf(\delta) > \nu = |[\nu]^{\sigma+1}|$, it follows that there exists some $z \in [\nu]^{\sigma+1}$ for which $\{\alpha \in A \mid z_{\alpha} = z\}$ is in J_{δ}^+ . As $E \subseteq E_z$, this is a contradiction. \dashv

Let z be given by the preceding claim. Then Φ^z is as sought.

Corollary 5.3. Suppose $\nu < \lambda < \kappa$ are infinite regular cardinals, and $S \subseteq E_{\lambda}^{\kappa}$.

If $\operatorname{CG}_{\lambda}(S, E_{\nu}^{\kappa})$ holds, then there exists a λ -bounded C-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ satisfying the following. For every club $D \subseteq \kappa$, for every $n < \omega$, there exists a $\delta \in S$ such that $\sup\{\beta < \delta \mid \operatorname{succ}_n(C_\delta \setminus \beta) \subseteq D\} = \delta$. \square

Theorem 5.4. Suppose $\xi \leq \kappa$ are uncountable cardinals. For every stationary $S \subseteq E_{>\omega_1}^{\kappa}$, $\operatorname{CG}_{\xi}(S, E_{\omega}^{\kappa}, 1, \vec{J})$ implies $\operatorname{CG}_{\xi}(S, \kappa, \omega, \vec{J})$;

Proof. Suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, E_{\omega}^{\kappa}, 1, \vec{J})$, with $S \subseteq E_{>\omega}^{\kappa}$. In particular, $\kappa \geq \aleph_2$. Fix an auxiliary C-sequence $\vec{e} = \langle e_{\gamma} \mid \gamma < \kappa \rangle$ such that $otp(e_{\gamma}) = cf(\gamma)$ for every $\gamma < \kappa$. In what follows we shall use the operator Φ_D from Definition 2.12.

Claim 5.4.1. There is a club $D \subseteq \kappa$ such that for every club $E \subseteq \kappa$ there is a $\delta \in S$ such that

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\omega}^{\kappa} \& \Phi_D(e_{\gamma}) \subseteq E\} \in J_{\delta}^+.$$

Proof. Suppose that the claim does not hold. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that for every $\delta \in S$,

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\omega}^{\kappa} \& \Phi_D(e_{\gamma}) \subseteq F^D\} \in J_{\delta}.$$

Let $\mu := \aleph_1$ so that $\mu < \kappa$. We construct now a \subseteq -decreasing sequence $\langle D_i \mid$ $i \leq \mu$ of clubs in κ as follows:

- $\begin{array}{ll} (\mathrm{i}) & D_0:=\kappa;\\ (\mathrm{ii}) & D_{i+1}:=D_i\cap F^{D_i}; \end{array}$
- (iii) for $i \in \operatorname{acc}(\mu+1)$, $D_i := \bigcap_{i' < i} D_{i'}$.

Since $\mu < \kappa$, all these are club in κ . Finally, consider the club

$$D^* := \{ \gamma < \kappa \mid \operatorname{otp}(D_\mu \cap \gamma) = \gamma \}$$

As $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, E_{\omega}^{\kappa}, 1, \vec{J})$, let us pick $\delta \in S$ such that

$$A := \{ \alpha < \delta \mid \min(C_{\delta} \setminus (\alpha + 1)) \in D^* \cap E_{\omega}^{\kappa} \}$$

is in J_{δ}^+ .

For every $i < \mu$, by the choice of F^{D_i} , the following set is in J_{δ} :

$$A_i := \{ \alpha < \delta \mid \gamma := \min(C_\delta \setminus (\alpha + 1)) \& \gamma \in E_\omega^\kappa \& \Phi_{D_i}(e_\gamma) \subseteq F^{D_i} \}.$$

As J_{δ} is $cf(\delta)$ -additive, and $cf(\delta) > \omega_1$, we may now fix $\alpha \in A \setminus \bigcup_{i < \mu} A_i$. Set $\gamma := \min(C_{\delta} \setminus (\alpha + 1))$, so that $\gamma \in D^* \cap E_{\omega}^{\kappa}$. Since $\gamma \in D^* \subseteq \operatorname{acc}(D_i)$ for every $i \leq \mu$, we have that $\Phi_{D_i}(e_{\gamma}) \subseteq D_i$ for every $i < \mu$.

For $\beta \in e_{\gamma}$, let $\beta_i := \sup(\beta \cap D_i)$. Since $\langle D_i \mid i \leq \mu \rangle$ is \subseteq -decreasing, $\langle \beta_i \mid$ $i < \mu$ is a non-increasing sequence, and hence it must stabilize beyond some ordinal $j(\beta) < \mu$. That is, for every $i \ge j(\beta)$ we have $\beta_i = \beta_{j(\beta)}$. Let $i^* := \sup_{\beta \in e_{\gamma}} j(\beta)$, and note that $i^* < \mu$, since $\mu = \aleph_1 = |e_{\gamma}|^+$. In particular, this implies that

$$\Phi_{D_{i^*}}(e_{\gamma}) = \Phi_{D_{i^*+1}}(e_{\gamma}) \subseteq D_{i^*+1} \subseteq D_{i^*} \cap F^{D_{i^*}}$$

On the other hand, since $\alpha \notin A_{i^*}$, we have that $\Phi_{D_{i^*}}(e_{\gamma}) \nsubseteq F^{D_{i^*}}$. This is a contradiction.

Let $D \subseteq \kappa$ be a club as given by the preceding claim. Consider a new *C*-sequence $\vec{C}^{\bullet} = \langle C^{\bullet}_{\delta} \mid \delta \in S \rangle$ defined via:

$$C^{\bullet}_{\delta} := C_{\delta} \cup \{ \Phi_D(e_{\gamma}) \setminus \sup(C_{\delta} \cap \gamma) \mid \gamma \in \operatorname{nacc}(C_{\delta}) \cap E^{\kappa}_{\omega} \}.$$

Note that for any $\delta \in S$, since $\operatorname{otp}(C_{\delta}) \leq \xi$ and for each $\gamma \in E_{\omega}^{\kappa}$ we have $\operatorname{otp}(\Phi_D(e_{\gamma})) \leq \operatorname{otp}(e) \leq \omega$, we have that the ordertype of every initial segment of C_{δ}^{\bullet} is strictly less than ξ , and hence $\operatorname{otp}(C_{\delta}^{\bullet}) \leq \xi$.

Now, if $E \subseteq \kappa$ is a club, then by the choice of D there is some $\delta \in S$ such that

$$A := \{ \alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\omega}^{\kappa} \& \Phi_D(e_{\gamma}) \subseteq \operatorname{acc}(D) \cap E \}$$

is in J_{δ}^+ . For every $\alpha \in A$, if we let $\gamma_{\alpha} := \min(C_{\delta}^{\bullet} \setminus (\alpha + 1))$, then $\gamma_{\alpha} \in E_{\omega}^{\kappa}$ and $C_{\delta}^{\bullet} \cap [\alpha, \gamma_{\alpha})$ is equal to $\Phi_D(e_{\gamma_{\alpha}}) \setminus \alpha$, which is an end segment of $\Phi_D(e_{\gamma_{\alpha}})$. Since any end segment of $\Phi_D(e_{\gamma_{\alpha}})$ has ordertype ω as well, it follows that for any $\alpha \in A$, there is an end segment of $C_{\delta}^{\bullet} \cap \gamma_{\alpha}$ of ordertype ω which is contained in E. Since this interval of ordertype ω which is contained in E also contains ω successive non-accumulation points of $C_{\delta}^{\bullet} \cap \gamma_{\alpha}$, we infer that

$$B := \{ \beta < \delta \mid \operatorname{succ}_{\omega}(C^{\bullet}_{\delta} \setminus \beta) \subseteq E \}$$

covers A. In particular, $B \in J^+_{\delta}$.

Theorem 5.5. Let $\sigma < \nu < \xi \leq \kappa$ be infinite cardinals. Suppose that $S \subseteq E_{>\nu}^{\kappa}$ and $T \subseteq \kappa$ are stationary sets.

- (1) If $\operatorname{CG}_{\xi}(S, E_{\sigma}^{\kappa} \cap \operatorname{Tr}(T), 1, \vec{J})$ holds, then so does $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$;
- (2) If $\operatorname{CG}_{\xi}(S, E_{\nu}^{\kappa} \cap \operatorname{Tr}(T), 1, \vec{J})$ holds, then so does $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$.

Proof. For the proof of both cases, we fix an auxiliary C-sequence $\vec{e} = \langle e_{\gamma} | \gamma < \kappa \rangle$ such that $\operatorname{otp}(e_{\gamma}) = \operatorname{cf}(\gamma)$ for every $\gamma < \kappa$.

(1) Suppose that $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, E_{\sigma}^{\kappa} \cap \operatorname{Tr}(T), 1, \vec{J})$. Let Φ^{B} be the operator from Definition 2.8.

Claim 5.5.1. There is a club $D \subseteq \kappa$ such that for every club $E \subseteq \kappa$ there is a $\delta \in S$ such that

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\sigma}^{\kappa} \cap \operatorname{Tr}(T) \& \Phi^{D \cap T}(e_{\gamma}) \subseteq E\} \in J_{\delta}^{+}.$$

Proof. Suppose that the claim does not hold. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that, for every $\delta \in S$,

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \gamma \in E_{\sigma}^{\kappa} \cap \operatorname{Tr}(T) \& \Phi^{D \cap T}(e_{\gamma}) \subseteq F^{D}\} \in J_{\delta}.$$

Set $\mu := \sigma^+$, so that $\mu \leq \nu < \kappa$. We construct now a \subseteq -decreasing sequence $\langle D_i | i \leq \mu \rangle$ of clubs in κ as follows:

(i) $D_0 := \kappa;$

(ii)
$$D_{i+1} := D_i \cap F^{D_i}$$

(iii) for $i \in \operatorname{acc}(\mu+1)$, $D_i := \bigcap_{i' < i} D_{i'}$.

Next, let $D^* := \operatorname{acc}(D_{\mu})$ and fix $\delta \in S$ such that

$$A := \{ \alpha < \delta \mid \min(C_{\delta} \setminus (\alpha + 1)) \in D^* \cap E_{\sigma}^{\kappa} \cap \operatorname{Tr}(T) \}$$

is in J_{δ}^+ .

For every $i < \mu$, as $D_{i+1} \subseteq F^{D_i}$, the following set

 $A_i := \{ \alpha < \delta \mid \gamma := \min(C_\delta \setminus (\alpha + 1)) \& \gamma \in E_\sigma^\kappa \cap \operatorname{Tr}(T) \& \Phi^{D_i \cap T}(e_\gamma) \subseteq D_{+i} \}$

is in J_{δ} . As $\operatorname{cf}(\delta) > \nu \ge \mu$, we may fix $\alpha \in A \setminus \bigcup_{i < \mu} A_i$. Set $\gamma := \min(C_{\delta} \setminus (\alpha + 1))$, so that $\gamma \in D^* \cap E_{\sigma}^{\kappa} \cap \operatorname{Tr}(T)$.

For every $i < \mu$, $\gamma \in \operatorname{acc}(D_i) \cap \operatorname{Tr}(T)$, so that $e_{\gamma} \cap D_i \cap T$ is stationary in γ , and hence $\Phi^{D_i \cap T}(e_{\gamma}) = \operatorname{cl}(e_{\gamma} \cap D_i \cap T)$. Thus, for every $i < \mu$, as $\alpha \notin A_i$, it must be the case that $\Phi^{D_i \cap T}(e_{\gamma}) \nsubseteq D_{i+i}$; but D_{i+1} is closed, so that, in fact, $e_{\gamma} \cap D_i \cap T \nsubseteq D_{i+i}$. For each $i < \mu$, pick $\beta_i \in (e_{\gamma} \cap D_i \cap T) \setminus D_{i+i}$. As $|e_{\gamma}| < \mu$, we may now fix $(i, j) \in [\mu]^2$ such that $\beta_i = \beta_j$. So $\beta_i \notin D_{i+1}$ while $\beta_j \in D_j \subseteq D_{i+1}$. This is a contradiction.

Let $D \subseteq \kappa$ be given by the preceding claim. Consider the *C*-sequence $\vec{C}^{\bullet} = \langle C_{\delta}^{\bullet} | \delta \in S \rangle$ defined via:

$$C^{\bullet}_{\delta} := C_{\delta} \cup \{ \Phi^{D \cap T}(e_{\gamma}) \setminus \sup(C_{\delta} \cap \gamma) \mid \gamma \in \operatorname{nacc}(C_{\delta}) \cap E^{\kappa}_{\sigma} \}$$

It is clear that \vec{C}^{\bullet} witnesses $\operatorname{CG}_{\xi}(S, T, \sigma, \vec{J})$.

(2) Suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, E_{\nu}^{\kappa} \cap \operatorname{Tr}(T), 1, \vec{J})$. For all $\gamma < \kappa$ and $\epsilon < \nu$, let $e_{\gamma}^{\epsilon} := \{\beta \in e_{\gamma} \cap T \mid \operatorname{otp}(e_{\gamma} \cap \beta) < \epsilon\}$ so that it is an initial segment of $e_{\gamma} \cap T$.

Claim 5.5.2. There is $\epsilon < \nu$ such that, for every club $E \subseteq \kappa$, there is $\delta \in S$ with

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \operatorname{otp}(e_{\gamma}^{\epsilon} \cap (E \setminus \alpha)) > \sigma\} \in J_{\delta}^{+}.$$

Proof. Otherwise, pick a counterexample E_{ϵ} for each $\epsilon < \nu$, and set $E := \bigcap_{\epsilon < \nu} E_{\epsilon}$. Pick $\delta \in S$ such that

$$A := \{ \alpha < \delta \mid \min(C_{\delta} \setminus (\alpha + 1)) \in \operatorname{acc}(E) \cap \operatorname{Tr}(T) \cap E_{\nu}^{\kappa} \}$$

is in J_{δ}^+ . For every $\alpha \in A$, if we let $\gamma_{\alpha} := \min(C_{\delta} \setminus (\alpha + 1))$, then $e_{\gamma} \cap E \cap T$ is a stationary subset of γ of ordertype ν , so there exists some $\epsilon_{\alpha} < \nu$ such that $\operatorname{otp}(e_{\gamma_{\alpha}}^{\epsilon_{\alpha}} \cap E \setminus \alpha) > \sigma$. As J_{δ} is $\operatorname{cf}(\delta)$ -additive and $\operatorname{cf}(\delta) > \nu$, there must exist some $\epsilon < \nu$ for which $A_{\epsilon} := \{\alpha \in A \mid \epsilon_{\alpha} = \epsilon\}$ is in J_{δ}^+ . But $E \subseteq E_{\epsilon}$. This is a contradiction.

Let ϵ be given by the claim.

Claim 5.5.3. There is a club $D \subseteq \kappa$ such that for every club $E \subseteq \kappa$ there is a $\delta \in S$ such that

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& e_{\gamma}^{\epsilon} \cap D \subseteq E \& \operatorname{otp}(e_{\gamma}^{\epsilon} \cap (D \setminus \alpha)) > \sigma\} \in J_{\delta}^{+}.$$

Proof. Suppose that the claim does not hold. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that, for every $\delta \in S$,

$$\{\alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& e_{\gamma}^{\epsilon} \cap D \subseteq F^{D} \& \operatorname{otp}(e_{\gamma}^{\epsilon} \cap (D \setminus \alpha)) > \sigma\} \in J_{\delta}.$$

We construct now a \subseteq -decreasing sequence $\langle D_i \mid i \leq \nu \rangle$ of clubs in κ as follows:

- (i) $D_0 := \kappa;$
- (ii) $D_{i+1} := D_i \cap F^{D_i};$
- (iii) for $i \in \operatorname{acc}(\nu+1)$, $D_i := \bigcap_{i' < i} D_{i'}$.
- Let $D^* := D_{\nu}$ and fix $\delta \in S$ such that

$$A := \{ \alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& \operatorname{otp}(e_{\gamma}^{\epsilon} \cap (D^* \setminus \alpha)) > \sigma \}$$

is in J^+_{δ} .

For every $i < \nu$, the following set

$$\begin{split} A_i &:= \{ \alpha < \delta \mid \gamma := \min(C_{\delta} \setminus (\alpha + 1)) \& e_{\gamma}^{\epsilon} \cap D_i \subseteq F^{D_i} \& \operatorname{otp}(e_{\gamma}^{\epsilon} \cap (D_i \setminus \alpha)) > \sigma \} \\ \text{is in } J_{\delta}. \text{ Fix } \alpha \in A \setminus \bigcup_{i < \nu} A_i. \text{ Set } \gamma := \min(C_{\delta} \setminus (\alpha + 1)), \text{ so that } \operatorname{otp}(e_{\gamma}^{\epsilon} \cap (D^* \setminus \alpha)) > \sigma. \end{split}$$

For every $i < \nu$, since $D^* \subseteq D_i$ and $D_{i+1} \subseteq F^{D_i}$, we have that $\operatorname{otp}(e_{\gamma}^{\epsilon} \cap (D_i \setminus \alpha)) > \sigma$, and hence it must be the case that $D_i \cap e_{\gamma}^{\epsilon} \nsubseteq F^{D_i}$, and therefore, $D_i \cap e_{\gamma}^{\epsilon} \nsubseteq D_{i+1}$. So $\langle D_i \cap e_{\gamma}^{\epsilon} \mid i < \nu \rangle$ is a strictly \subseteq -decreasing sequence of subsets of e_{γ}^{ϵ} , contradicting the fact that $|e_{\gamma}^{\epsilon}| \le |\epsilon| < \nu$.

Let *D* be given by the claim. As $\epsilon < \nu$, for every $\gamma < \kappa$, $|e_{\gamma}^{\epsilon}| < \nu$. It altogether follows that the *C*-sequence $\vec{C}^{\bullet} = \langle C_{\delta}^{\bullet} | \delta \in S \rangle$ defined via:

$$C^{\bullet}_{\delta} := C_{\delta} \cup \{ \operatorname{cl}(D \cap e^{\epsilon}_{\gamma}) \setminus \sup(C_{\delta} \cap \gamma) \mid \gamma \in \operatorname{nacc}(C_{\delta}) \},$$

is as sought.

Corollary 5.6. Let $\mu < \sigma < \sigma^+ < \lambda < \kappa$ be infinite regular cardinals. Then $\operatorname{CG}_{\lambda}(E_{\lambda}^{\kappa}, E_{\sigma}^{\kappa}, 1, \vec{J})$ implies $\operatorname{CG}_{\lambda}(E_{\lambda}^{\kappa}, E_{\mu}^{\kappa}, \sigma, \vec{J})$.

Proof. Appeal to Theorem 5.5(1) with $\nu := \sigma^+, \xi := \lambda, S := E_{\lambda}^{\kappa}$ and $T := E_{\mu}^{\kappa}$. \Box

Corollary 5.7. For every successor cardinal λ , if $CG_{\xi}(E_{\lambda}^{\lambda^{+}}, E_{<\lambda}^{\lambda^{+}}, 1)$ holds, then so does $CG_{\xi}(E_{\lambda}^{\lambda^{+}}, \lambda^{+}, 2)$.

Proof. By Theorem 5.2, using $\sigma := 2, \lambda := \nu^+, \kappa := \lambda^+, S := E_{\lambda}^{\kappa}$, and $\vec{J} := \langle J^{\mathrm{bd}}[\delta] \mid \delta \in S \rangle$.

Remark 5.8. This shows that Clause (4) of [Asp14, Theorem 1.6] follows from Clause (5) of the same theorem, provided that the cardinal κ there is a successor cardinal.

6. MOVING BETWEEN IDEALS

As shown in the Section 4, it is easier to partition a witness for $\operatorname{CG}_{\xi}(S, T, \sigma, \overline{J})$ in the case that the ideals in \overline{J} are normal. So, we address here the problem of deriving $\operatorname{CG}_{\xi}(S, T, \sigma, \langle \operatorname{NS}_{\delta} | \delta \in S \rangle)$ from $\operatorname{CG}_{\xi}(S, T, \sigma, \langle J^{\operatorname{bd}}[\delta] | \delta \in S \rangle)$. The key lemma is Lemma 6.3. In Theorem 6.4 it is used to improve results from Section 4. At successor cardinals, Lemma 6.3 is particularly useful as seen by the main result of this section, which combines Theorems A and E:

Corollary 6.1. Suppose that λ is a successor cardinal, and $S \subseteq E_{\lambda}^{\lambda^+}$ is stationary. *Then:*

- (1) $\operatorname{CG}_{\lambda}(S, E_{\lambda}^{\lambda^{+}}, 1, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$ holds;
- (2) $\operatorname{CG}_{\lambda}(S, E_{<\lambda}^{\lambda^+})$ implies $\operatorname{CG}_{\lambda}(S, \lambda^+, n, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$ for every $n < \omega$;
- (3) If $\lambda > \aleph_1$, then $\operatorname{CG}_{\lambda}(S, E_{<\lambda}^{\lambda^+})$ implies $\operatorname{CG}_{\lambda}(S, \lambda^+, \omega, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$.

Proof. Let ν denote the predecessor of λ .

(1) By Fact 3.25 and Theorem 6.11(1) below.

(2) Suppose that $\operatorname{CG}_{\lambda}(S, E_{<\lambda}^{\lambda^+})$, equivalently $\operatorname{CG}_{\lambda}(S, E_{\geq\omega}^{\lambda^+} \cap E_{\leq\nu}^{\lambda^+})$, holds. Then, by Corollary 5.1, also $\operatorname{CG}_{\lambda}(S, \lambda^+, n)$ holds for every $n < \omega$. Finally, By Theorem 6.11(1) below, moreover $\operatorname{CG}_{\lambda}(S, \lambda^+, n, \langle \operatorname{NS}_{\delta} | \delta \in E_{\lambda}^{\lambda^+} \rangle)$ holds for every $n < \omega$.

(3) Assuming that ν is uncountable, by Corollary 5.1, $\operatorname{CG}_{\lambda}(S, E_{\geq \omega}^{\lambda^+} \cap E_{\leq \nu}^{\lambda^+})$ implies $\operatorname{CG}_{\lambda}(S, \lambda^+, \omega)$, which, by Theorem 6.11(1) below, implies $\operatorname{CG}_{\lambda}(S, \lambda^+, \omega, \langle \operatorname{NS}_{\delta} | \delta \in E_{\lambda}^{\lambda^+} \rangle)$.

Lemma 6.2. Let $\aleph_0 < \xi < \kappa$ and $S \subseteq E_{\xi}^{\kappa}$ be stationary. Assume $1 \leq \sigma < \xi$.

If $\vec{J} = \langle J_{\delta} | \delta \in S \rangle$ is such that $NS_{\delta} \subseteq J_{\delta}$ for all δ , then $CG(S, T, \sigma, \vec{J})$ implies $CG_{\xi}(S, T, \sigma, \vec{J})$.

Proof. Let $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ be a witness to $CG(S, T, \sigma, \vec{J})$. For each $\delta \in S$, define a function $f_{\delta} : \delta \to \delta$ via

$$f_{\delta}(\beta) := \sup(\operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta)) + 1,$$

then fix a club e_{δ} in δ of ordertype $cf(\delta)$ consisting of closure points of f_{δ} , and finally let C^{\bullet}_{δ} be the ordinal closure below δ of the following set:

$$\bigcup \{ \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \mid \beta \in e_{\delta} \}$$

To see that $\langle C^{\bullet}_{\delta} | \delta \in S \rangle$ witnesses $CG_{\xi}(S, T, \sigma, \vec{J})$, let D be a club in κ . Pick $\delta \in S$ for which the following set is in J^+_{δ} :

$$B := \{ \beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \}.$$

Then $e_{\delta} \cap B \in J_{\delta}^+$, and for every $\beta \in e_{\delta} \cap B$, $\operatorname{succ}_{\sigma}(C_{\delta}^{\bullet} \setminus \beta) = \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta)$. \Box

Lemma 6.3. Suppose that $\vec{C} = \langle C_{\delta} | \delta \in S \rangle$ witnesses $\operatorname{CG}(S, T, \sigma)$, with $S \subseteq E_{>\omega}^{\kappa}$ and $T \subseteq \kappa$. Then there exists a *C*-sequence $\vec{e} = \langle e_{\delta} | \delta \in S \rangle$ such that, for every club $D \subseteq \kappa$, there exists $\delta \in S$ such that the following set is stationary in δ :

$$\{\alpha \in e_{\delta} \mid \exists \beta \in C_{\delta} \left[\alpha \leq \beta < \min(e_{\delta} \setminus (\alpha + 1)) \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \right] \}.$$

Proof. Suppose not. For every $\delta \in S$, let $e_{\delta}^{0} := C_{\delta}$. Next, suppose that $i < \omega$ and that $\langle e_{\delta}^{i} \mid \delta \in S \rangle$ has already been defined. By assumption, we can find a club $D_{i} \subseteq \kappa$ such that, for every $\delta \in S$, the following set is nonstationary in δ :

$$\{\alpha \in e^i_{\delta} \mid \exists \beta \in C_{\delta} \left[\alpha \leq \beta < \min(e^i_{\delta} \setminus (\alpha + 1)) \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D_i \cap T \right] \},\$$

so let us pick a subclub e_{δ}^{i+1} of e_{δ}^{i} disjoint from it.

Put $D := \bigcap_{i < \omega} D_i$. By the choice of \vec{C} , let us now pick $\delta \in S$ such that

$$\sup\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} = \delta.$$

Pick $\beta < \delta$ above $\min(\bigcap_{i < \omega} e^i_{\delta})$ such that $\operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T$. Consider the ordinal

$$\gamma := \min(\operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta)),$$

and then, for every $i < \omega$, let $\alpha_i := \sup(e^i_{\delta} \cap \gamma)$. As $\operatorname{acc}(e^i_{\delta}) \subseteq \operatorname{acc}(C_{\delta})$, and γ is in $\operatorname{nacc}(C_{\delta})$ and above $\min(e^i_{\delta})$, we infer that $\alpha_i \in e^i_{\delta} \cap \gamma$. As $\langle e^i_{\delta} \mid i < \omega \rangle$ is a \subseteq -decreasing chain, $\langle \alpha_i \mid i < \omega \rangle$ is \leq -decreasing, so we may find a large enough $i < \omega$ such that $\alpha_{i+1} = \alpha_i$. In particular, $\alpha_i \in e^{i+1}_{\delta}$, so by the choice of e^{i+1}_{δ} ,

$$\forall \beta \in C_{\delta} \left[\alpha \leq \beta < \min(e_{\delta}^{i} \setminus (\alpha + 1)) \rightarrow \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \notin D_{i} \cap T \right].$$

On the other hand, since $\alpha_i = \sup(e_{\delta}^i \cap \gamma)$, it is the case that $\min(e_{\delta}^i \setminus (\alpha_i + 1)) \ge \gamma$. Recalling also that $e_{\delta}^0 \subseteq C_{\delta}$, altogether $\alpha_i \le \beta < \min(e_{\delta}^i \setminus (\alpha_i + 1))$, and

$$\operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \subseteq D_i \cap T.$$

This is a contradiction.

An immediate consequence of the preceding lemma is an improvement of Clauses (2) and (3) of Theorem 4.9, for the special case of $\vec{J} = \langle J^{\text{bd}}[\delta] \mid \delta \in S \rangle$.

Theorem 6.4. Suppose that \vec{C} witnesses $CG(S,T,\sigma)$ with $S \subseteq E_{\lambda}^{\kappa}$.

- (1) If onto(NS_{λ}, λ) holds, then $\lambda \in \Theta_2(\vec{C}, T, \sigma)$;
- (2) If unbounded(NS_{λ}, θ) holds and $\theta < \lambda$, then $\theta \in \Theta_2(\vec{C}, T, \sigma)$.

Proof. Let $\vec{e} = \langle e_{\delta} | \delta \in S \rangle$ be the corresponding *C*-sequence given by Lemma 6.3. Without loss of generality, $\operatorname{otp}(e_{\delta}) = \lambda$ for all $\delta \in S$. Define $\psi_{\delta} : \delta \to \lambda$ via $\psi_{\delta}(\beta) := \operatorname{otp}(e_{\delta} \cap \beta)$, so that, for every $A \subseteq \delta$, $\psi_{\delta}[A]$ is stationary in λ iff *A* is stationary in δ .

(1): Suppose that $onto(NS_{\delta}, \lambda)$ holds, and fix a colouring $c : [\lambda]^2 \to \lambda$ as in Theorem 4.10(2). As $nacc(\kappa)$ is in NS_{λ} , we may assume that for all $\eta < \alpha < \kappa$, $c(\eta, \alpha + 1) = c(\eta, \alpha)$. Now, a proof nearly identical to that of Claim 4.9.1 yields an $\eta < \lambda$ such that, for every club $D \subseteq \kappa$, there exists a $\delta \in S$, such that, for every $\tau < \lambda$:

 $\sup\{\beta < \delta \mid \eta < \psi_{\delta}(\beta) \& c(\eta, \psi_{\delta}(\beta)) = \tau \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} = \delta.$

Choose $\vec{h} = \langle h_{\delta} : \delta \to \lambda \mid \delta \in S \rangle$ satisfying $h_{\delta}(\beta) = c(\eta, \psi_{\delta}(\beta))$ for every $\delta \in S$ and $\beta < \delta$ such that $\eta < \psi_{\delta}(\beta)$. Then \vec{h} witnesses that $\lambda \in \Theta_2(\vec{C}, T, \sigma)$.

(2) Suppose that $\mathsf{unbounded}(J^{\mathrm{bd}}[\lambda], \theta)$ holds with $\theta < \lambda$, and fix a colouring $c : [\lambda]^2 \to \theta$ as in Theorem 4.10(1). For every club $D \subseteq \kappa$, for all $\delta \in S$ and $\eta < \lambda$, let $D(\eta, \delta)$ denote the set:

 $\{\tau < \theta \mid \sup\{\beta < \delta \mid \eta < \psi_{\delta}(\beta) \& c(\eta, \psi_{\delta}(\beta)) = \tau \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T\} = \delta\}.$

A proof nearly identical to that of Claim 4.9.2 yields an $\eta < \lambda$ such that, for every club $D \subseteq \kappa$, there exists $\delta \in S$, such that $|D(\eta, \delta)| = \theta$. The rest of the proof is now identical to that of Theorem 4.9(3).

Corollary 6.5. Suppose that \vec{C} witnesses $CG(S,T,\sigma)$ with $S \subseteq E_{\lambda}^{\kappa}$. If λ is not ineffable, then $\omega \in \Theta_2(\vec{C},T,\sigma)$.

Proof. By Theorem 6.4 and Corollary 4.19(3).

Motivated by Fact 4.16(1), we ask:

Question 6.6. Suppose that CG(S,T) holds for stationary $S \subseteq E_{>\omega}^{\kappa}$ and $T \subseteq \kappa$. Does there exist a cardinal $\mu < \kappa$ such that $CG(S,T,1, \langle NS_{\delta} | E_{\mu}^{\delta} | \delta \in S \rangle)$ holds?

Lemma 6.3 suggests the following variation of Definition 2.2.

Definition 6.7. $\operatorname{CG}_{\xi}(S, T, \frac{1}{2}, \vec{J})$ asserts the existence of a ξ -bounded *C*-sequence $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ such that, for every club $D \subseteq \kappa$ there is a $\delta \in S$ such that

$$\{\beta < \delta \mid (\beta, \min(C_{\delta} \setminus (\beta + 1))] \cap D \cap T \neq \emptyset\} \in J_{\delta}^+.$$

Corollary 6.8. For all stationary $S \subseteq E_{>\omega}^{\kappa}$ and $T \subseteq \kappa$, if CG(S,T) holds, then so does $CG(S,T, \frac{1}{2}, \langle NS_{\delta} | \delta \in S \rangle)$.

We now show that it is possible to upgrade $\sigma = \frac{1}{2}$ to $\sigma = 1$, but at the cost of losing control over the set T.

Lemma 6.9. Suppose that $S \subseteq E_{>\omega}^{\kappa}$ is stationary.

For every C-sequence $\langle C_{\delta} | \delta \in S \rangle$ witnessing $\operatorname{CG}(S, \kappa, \frac{1}{2}, \overline{J})$, there exists a postprocessing^{*} function $\Phi : \mathcal{K}(\kappa) \to \mathcal{K}(\kappa)$ such that $\langle \Phi(C_{\delta}) | \delta \in S \rangle$ witnesses $\operatorname{CG}(S, \kappa, 1, \overline{J})$.

Proof. We shall make use of the operator Φ_D from Definition 2.12.

Claim 6.9.1. There exists a club $D \subseteq \kappa$ such that, $\langle \Phi_D(C_\delta) | \delta \in S \rangle$ witnesses $CG(S, \kappa, 1, \vec{J})$.

Proof. Suppose not. In this case, for every club $D \subseteq \kappa$, there is a club $F^D \subseteq \kappa$ such that for every $\delta \in S$

$$\{\beta < \delta \mid \min(\Phi_D(C_\delta) \setminus (\beta + 1)) \in F^D\} \in J_\delta.$$

Construct a \subseteq -decreasing sequence $\langle D_n | n < \omega \rangle$ of clubs in κ by letting $D_0 := \kappa$ and $D_{n+1} := \operatorname{acc}(D_n) \cap F^{D_n}$ for every $n < \omega$. Set $D := \bigcap_{n < \omega} D_n$ and then pick $\delta \in S$ for which the following set is in J_{δ}^+ :

$$B := \{ \beta \in C_{\delta} \mid (\beta, \min(C_{\delta} \setminus (\beta + 1))] \cap D \neq \emptyset \}.$$

In particular, $\delta \in \operatorname{acc}(D)$, so that, for every $n < \omega$,

$$\Phi_{D_n}(C_{\delta}) = \{ \sup(D_n \cap \eta) \mid \eta \in C_{\delta}, \eta > \min(D_n) \}.$$

Now, as $cf(\delta) > \omega$ and J_{δ} is $cf(\delta)$ -complete, the following set is nonempty:

$$B \setminus \bigcup_{n < \omega} \{\beta < \delta \mid \min(\Phi_{D_n}(C_\delta) \setminus (\beta + 1)) \in F^{D_n}\},\$$

so we may pick in it some ordinal β . Set $\gamma := \min(C_{\delta} \setminus (\beta + 1))$. As $\beta \in B$, we know that $D \cap (\beta, \gamma] \neq \emptyset$. In particular, for every $n < \omega$, $\operatorname{acc}(D_n) \cap (\beta, \gamma] \neq \emptyset$ and $\beta_n := \sup(D_n \cap \gamma)$ is an element of D_n greater than β , so that

$$\min(\Phi_{D_n}(C_{\delta}) \setminus (\beta + 1)) = \{\beta_n\}$$

As $\langle D_n \mid n < \omega \rangle$ is a \subseteq -decreasing chain, we may fix $n < \omega$ such that $\beta_{n+1} = \beta_n$. Then $\min(\Phi_{D_n}(C_{\delta}) \setminus (\beta+1)) = \beta_n = \beta_{n+1} \in D_{n+1} \subseteq F^{D_n}$, contradicting the choice of β .

Let D be given by the preceding claim. Then $\Phi := \Phi_D$ is as sought.

Corollary 6.10. Suppose that $S \subseteq E_{>\omega}^{\kappa}$ is stationary. If $CG(S, \kappa)$ holds, then so does $CG(S, \kappa, 1, \langle NS_{\delta} | \delta \in S \rangle)$.

Proof. By Corollary 6.8 and Lemma 6.9.

Note that the preceding result is restricted to $\sigma := 1$ and $T := \kappa$. We now provide a condition sufficient for waiving this restriction.

Theorem 6.11. Suppose that ξ is an infinite successor cardinal, and $S \subseteq E_{\xi}^{\kappa}$ is stationary.

(1) If $\operatorname{CG}_{\xi}(S,T,\sigma)$ holds, then so does $\operatorname{CG}_{\xi}(S,T,\sigma,\langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$; (2) If $\operatorname{CG}_{\xi}(S,T,\frac{1}{2})$ holds, then so does $\operatorname{CG}_{\xi}(S,T,1,\langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$. *Proof.* We provide a proof of Clause (1) and leave the modification of the argument to obtain Clause (2) to the reader.

As $\operatorname{CG}_{\xi}(S, T, \xi)$ is equivalent to $\operatorname{CG}(S, T, \kappa)$, by Lemma 2.10, we may assume that $\sigma < \xi$. Now, suppose that $\vec{C} = \langle C_{\delta} \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, T, \sigma)$, and let $\vec{e} = \langle e_{\delta} \mid \delta \in S \rangle$ be the corresponding *C*-sequence given by Lemma 6.3. For each $\delta \in S$, by possibly shrinking e_{δ} as in the proof of Lemma 6.2, we may assume that for every $\gamma \in e_{\delta}$ and every $\beta < \gamma$, $\sup(\operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta)) < \gamma$.

Let μ be such that $\xi = \mu^+$, and then, for all $\delta \in S$ and $\alpha \in e_{\delta}$, let $\varphi_{\delta,\alpha}$ be some surjection from μ to $C_{\delta} \cap [\alpha, \min(e_{\delta} \setminus (\alpha + 1)))$.

For every $i < \mu$, let C^i_{δ} be the ordinal closure below δ of the following set:

$$\bigcup \{ \operatorname{succ}_{\sigma}(C_{\delta} \setminus \varphi_{\delta,\alpha}(i)) \mid \alpha \in e_{\delta} \}.$$

Claim 6.11.1. There exists $i < \mu$, such that, for every club $D \subseteq \kappa$, there exists $\delta \in S$ for which the following set is stationary in δ :

$$\{\beta < \delta \mid \operatorname{succ}_{\sigma}(C^i_{\delta} \setminus \beta) \subseteq D \cap T\}.$$

Proof. Suppose not. For each $i < \mu$, pick a counterexample club $D_i \subseteq \kappa$. Consider the club $D := \bigcap_{i < \mu} D_i$. By the choice of \vec{e} , pick $\delta \in S$ such that

$$A := \{ \alpha \in e_{\delta} \mid \exists \beta \in C_{\delta} \left[\alpha \leq \beta < \min(e_{\delta} \setminus (\alpha + 1)) \& \operatorname{succ}_{\sigma}(C_{\delta} \setminus \beta) \subseteq D \cap T \right] \}$$

is stationary. For every $\alpha \in A$, pick $i_{\alpha} < \mu$ such that $\beta_{\alpha} := \varphi_{\delta,\alpha}(i)$ witnesses that $\alpha \in A$, that is, such that $\operatorname{succ}_{\sigma}(C_{\delta} \setminus \varphi_{\delta,\alpha}(i)) \subseteq D \cap T$. As $\operatorname{cf}(\delta) = \xi > \mu$, there must exist some stationary $A^* \subseteq A$ on which the map $\alpha \mapsto i_{\alpha}$ is constant, with value, say, i^* . For every pair $\alpha < \alpha'$ of ordinals from e_{δ} ,

$$\sup(\operatorname{succ}_{\sigma}(C_{\delta} \setminus \varphi_{\delta,\alpha}(i^*))) < \alpha' \leq \min(\operatorname{succ}_{\sigma}(C_{\delta} \setminus \varphi_{\delta,\alpha'}(i^*))),$$

so, recalling the definition of C_{δ}^{i*} , for every $\alpha \in A^*$,

$$\operatorname{succ}_{\sigma}(C_{\delta}^{i^*} \setminus \alpha) = \operatorname{succ}_{\sigma}(C_{\delta} \setminus \varphi_{\delta,\alpha}(i^*)) \subseteq D \cap T,$$

contradicting the fact that $D \subseteq D_{i^*}$.

Let $i < \mu$ be given by the preceding. Then $\langle C_{\delta}^i \mid \delta \in S \rangle$ witnesses $\operatorname{CG}_{\xi}(S, T, \sigma, \langle \operatorname{NS}_{\delta} \mid \delta \in S \rangle)$.

Remark 6.12. The above ordertype restriction cannot be waived, that is, the hypothesis $CG_{\xi}(S, T, \sigma)$ in Theorem 6.11 cannot be relaxed to $CG(S, T, \sigma)$.

By Theorem 3.27(1), if there exists a nonreflecting stationary subset of $E_{\aleph_0}^{\aleph_2}$, then $\operatorname{CG}(E_{\aleph_1}^{\aleph_2}, E_{\aleph_0}^{\aleph_2})$ holds, and then, by Theorem 5.2, using $\xi = \kappa = \aleph_2$ and $S = E_{\aleph_1}^{\aleph_2}$, so does $\operatorname{CG}(E_{\aleph_1}^{\aleph_2}, \omega_2, 2)$. Now, if the ordertype restriction in Theorem 6.11 could have been waived, then this would imply that $\operatorname{CG}(E_{\aleph_1}^{\aleph_2}, \omega_2, 2, \langle \operatorname{NS}_{\delta} \mid \delta \in E_{\aleph_1}^{\aleph_2} \rangle)$ holds. In particular, by Lemma 6.2, $\operatorname{CG}_{\omega_1}(E_{\aleph_1}^{\aleph_2}, \omega_2, 2)$ holds. However, running the forcing from [Asp14, Theorem 1.6] over a model of \Box_{ω_1} , one gets a generic extension with a nonreflecting stationary subset of $E_{\aleph_0}^{\aleph_2}$ in which $\operatorname{CG}_{\omega_1}(E_{\aleph_1}^{\aleph_2}, \omega_2, 2)$ fails. This is a contradiction.

Remark 6.13. An obvious complication of the proof of Theorem 6.11 shows that if ξ an infinite successor cardinal, $S \subseteq E_{\xi}^{\kappa}$ is stationary, and \vec{C} is a ξ -bounded *C*-sequence over *S* such that $\theta \in \Theta_2(\vec{C}, T, \sigma)$, then there exists a ξ -bounded *C*sequence \vec{C}^{\bullet} over *S* such that $\theta \in \Theta_2(\vec{C}^{\bullet}, T, \sigma, \langle NS_{\delta} | \delta \in S \rangle)$.

Acknowledgments

The first author is supported by the Israel Science Foundation (grant agreement 2066/18). The second author is partially supported by the Israel Science Foundation (grant agreement 2066/18) and by the European Research Council (grant agreement ERC-2018-StG 802756).

References

- [AFMS13] David Asperó, Sy-David Friedman, Miguel Angel Mota, and Marcin Sabok. Baumgartner's conjecture and bounded forcing axioms. Ann. Pure Appl. Logic, 164(12):1178– 1186, 2013.
- [AK20] David Asperó and John Krueger. Parametrized measuring and club guessing. Fund. Math., 249(2):169–183, 2020.
- [AM10] Uri Abraham and Menachem Magidor. Cardinal arithmetic. In Handbook of set theory. Vols. 1, 2, 3, pages 1149–1227. Springer, Dordrecht, 2010.
- [Asp14] David Asperó. The consistency of a club-guessing failure at the successor of a regular cardinal. In *Infinity, computability, and metamathematics*, volume 23 of *Tributes*, pages 5–27. Coll. Publ., London, 2014.
- [BGKT78] S. Broverman, J. Ginsburg, K. Kunen, and F. D. Tall. Topologies determined by σideals on ω₁. Canadian J. Math., 30(6):1306–1312, 1978.
- [BHZ07] Andreas Blass, Tapani Hyttinen, and Yi Zhang. Mad families and their neighbors, 2007.
- [BLS17] Tomek Bartoszyński, Paul Larson, and Saharon Shelah. Closed sets which consistently have few translations covering the line. *Fund. Math.*, 237(2):101–125, 2017.
- [BR17a] Ari Meir Brodsky and Assaf Rinot. A microscopic approach to Souslin-tree constructions. Part I. Ann. Pure Appl. Logic, 168(11):1949–2007, 2017.
- [BR17b] Ari Meir Brodsky and Assaf Rinot. Reduced powers of Souslin trees. Forum Math. Sigma, 5:Paper No. e2, 82, 2017.
- [BR19a] Ari Meir Brodsky and Assaf Rinot. Distributive Aronszajn trees. Fund. Math., 245(3):217–291, 2019.
- [BR19b] Ari Meir Brodsky and Assaf Rinot. A remark on Schimmerling's question. Order, 36(3):525–561, 2019.
- [BR21] Ari Meir Brodsky and Assaf Rinot. A microscopic approach to Souslin-tree constructions. Part II. Ann. Pure Appl. Logic, 172(5):102904, 2021.
- [Dza05] Mirna Dzamonja. Club guessing and the universal models. Notre Dame J. Formal Logic, 46(3):283–300, 2005.
- [Eis10a] Todd Eisworth. Club-guessing, stationary reflection, and coloring theorems. Ann. Pure Appl. Logic, 161(10):1216–1243, 2010.
- [Eis10b] Todd Eisworth. Successors of singular cardinals. In Handbook of set theory. Vols. 1, 2, 3, pages 1229–1350. Springer, Dordrecht, 2010.
- [EN09] Todd Eisworth and Peter Nyikos. Antidiamond principles and topological applications. Trans. Amer. Math. Soc., 361(11):5695–5719, 2009.
- [ES05] Todd Eisworth and Saharon Shelah. Successors of singular cardinals and coloring theorems. I. Arch. Math. Logic, 44(5):597–618, 2005.
- [ES09] Todd Eisworth and Saharon Shelah. Successors of singular cardinals and coloring theorems. II. J. Symbolic Logic, 74(4):1287–1309, 2009.
- [FK05] Matthew Foreman and Peter Komjath. The club guessing ideal: commentary on a theorem of Gitik and Shelah. J. Math. Log., 5(1):99–147, 2005.
- [FT05] Matthew Foreman and Stevo Todorcevic. A new Löwenheim-Skolem theorem. Trans. Amer. Math. Soc., 357(5):1693–1715, 2005.
- [GHZ21] Osvaldo Guzmán, Michael Hrušák, and Jindřich Zapletal. Preservation theorems for Namba forcing. Ann. Pure Appl. Logic, 172(2):Paper No. 102869, 12, 2021.
- [GS97] Moti Gitik and Saharon Shelah. Less saturated ideals. Proc. Amer. Math. Soc., 125(5):1523–1530, 1997.
- [GS22] Mohammad Golshani and Saharon Shelah. The measuring principle and the continuum hypothesis. 2022. arXiv: 2207.08048.

- [HHS09] F. Hernández-Hernández and Paul J. Szeptycki. A small Dowker space from a clubguessing principle. *Topology Proc.*, 34:351–363, 2009.
- [Hir07] James Hirschorn. Random trees under CH. Israel J. Math., 157:123-153, 2007.
- [Hof13] Douglas J. Hoffman. A Coloring Theorem for Inaccessible Cardinals. ProQuest LLC, Ann Arbor, MI, 2013. Thesis (Ph.D.)–Ohio University.
- [IL12a] Tetsuya Ishiu and Paul B. Larson. P_{max} variations for separating club guessing principles. J. Symbolic Logic, 77(2):532–544, 2012.
- [IL12b] Tetsuya Ishiu and Paul B. Larson. Some results about (+) proved by iterated forcing. J. Symbolic Logic, 77(2):515–531, 2012.
- [IR22a] Tanmay Inamdar and Assaf Rinot. Was Ulam right? I: Basic theory and subnormal ideals. *Topology Appl.*, 2022. Accepted December 2021.
- [IR22b] Tanmay Inamdar and Assaf Rinot. Was Ulam right? II: Small width and general ideals. http://assafrinot.com/paper/53, 2022. Submitted March 2022.
- [Ish05] Tetsuya Ishiu. Club guessing sequences and filters. J. Symbolic Logic, 70(4):1037–1071, 2005.
- [Ish15] Tetsuya Ishiu. The comparison of various club guessing principles. Ann. Pure Appl. Logic, 166(5):583–600, 2015.
- [Jen68] R. B. Jensen. Souslin's hypothesis is incompatible with V = L. Notices Amer. Math. Soc., 15(6):935, 1968.
- [Jen72] R. Björn Jensen. The fine structure of the constructible hierarchy. Ann. Math. Logic, 4:229–308; erratum, ibid. 4 (1972), 443, 1972. With a section by Jack Silver.
- [Koj95] Menachem Kojman. The A, B, C of pcf: a companion to pcf theory, part I. arXiv preprint math/9512201, 1995.
- [KS92] Menachem Kojman and Saharon Shelah. Nonexistence of universal orders in many cardinals. J. Symbolic Logic, 57(3):875–891, 1992.
- [KS93] Menachem Kojman and Saharon Shelah. μ -complete Souslin trees on μ^+ . Arch. Math. Logic, 32(3):195–201, 1993.
- [KS01] Menachem Kojman and Saharon Shelah. Fallen cardinals. Ann. Pure Appl. Logic, 109(1-2):117–129, 2001.
- [LH20] Chris Lambie-Hanson. On the growth rate of chromatic numbers of finite subgraphs. Adv. Math., 369:107176, 13, 2020.
- [LHR18] Chris Lambie-Hanson and Assaf Rinot. Knaster and friends I: closed colorings and precalibers. Algebra Universalis, 79(4):Art. 90, 39, 2018.
- [LHR19] Chris Lambie-Hanson and Assaf Rinot. Reflection on the coloring and chromatic numbers. Combinatorica, 39(1):165–214, 2019.
- [Mat09] Pierre Matet. Game ideals. Ann. Pure Appl. Logic, 158(1-2):23-39, 2009.
- [Moo08] Justin Tatch Moore. Aronszajn lines and the club filter. J. Symbolic Logic, 73(3):1029– 1035, 2008.
- [Nee14] Itay Neeman. Forcing with sequences of models of two types. Notre Dame J. Form. Log., 55(2):265–298, 2014.
- [Ost76] A. J. Ostaszewski. On countably compact, perfectly normal spaces. J. London Math. Soc. (2), 14(3):505–516, 1976.
- [RaS96] Andrzej Rosł anowski and Saharon Shelah. More forcing notions imply diamond. Arch. Math. Logic, 35(5-6):299–313, 1996.
- [Rin10a] Assaf Rinot. A relative of the approachability ideal, diamond and non-saturation. J. Symbolic Logic, 75(3):1035–1065, 2010.
- [Rin10b] Assaf Rinot. The search for diamonds. B. Symbolic Logic, 16(3):420–423, 2010.
- [Rin11] Assaf Rinot. Jensen's diamond principle and its relatives. In Set theory and its applications, volume 533 of Contemp. Math., pages 125–156. Amer. Math. Soc., Providence, RI, 2011.
- [Rin14a] Assaf Rinot. Chain conditions of products, and weakly compact cardinals. Bull. Symb. Log., 20(3):293–314, 2014.
- [Rin14b] Assaf Rinot. Complicated colorings. Math. Res. Lett., 21(6):1367–1388, 2014.
- [Rin14c] Assaf Rinot. The Ostaszewski square and homogeneous Souslin trees. Israel J. Math., 199(2):975–1012, 2014.
- [Rin15a] Assaf Rinot. Chromatic numbers of graphs large gaps. Combinatorica, 35(2):215–233, 2015.

- [Rin15b] Assaf Rinot. Putting a diamond inside the square. Bull. Lond. Math. Soc., 47(3):436– 442, 2015.
- [Rin17] Assaf Rinot. Higher Souslin trees and the GCH, revisited. Adv. Math., 311(C):510–531, 2017.
- [Rin19] Assaf Rinot. Souslin trees at successors of regular cardinals. MLQ Math. Log. Q., 65(2):200–204, 2019.
- [Rin22] Assaf Rinot. On the ideal $J[\kappa]$. Ann. Pure Appl. Logic, 173(2):Paper No. 103055, 13, 2022.
- [RZ21] Assaf Rinot and Jing Zhang. Transformations of the transfinite plane. Forum Math. Sigma, 9(e16):1–25, 2021.

[RZ22] Assaf Rinot and Jing Zhang. Complicated colorings, revisited. http://assafrinot.com/paper/52, 2022. Submitted February 2022.

[She94a] Saharon Shelah. Cardinal Arithmetic. In Cardinal Arithmetic, volume 29 of Oxford Logic Guides, chapter IX. Oxford University Press, 1994. Ch. IX of [She94b].

[She94b] Saharon Shelah. Cardinal arithmetic, volume 29 of Oxford Logic Guides. The Clarendon Press, Oxford University Press, New York, 1994. Oxford Science Publications.

[She94c] Saharon Shelah. There are Jonsson algebras in many inaccessible cardinals. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.

- [She97] Saharon Shelah. Colouring and non-productivity of ℵ₂-c.c. Ann. Pure Appl. Logic, 84(2):153–174, 1997.
- [She98] Saharon Shelah. *Proper and improper forcing*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 2nd edition, 1998.
- [She00] Saharon Shelah. On what I do not understand (and have something to say). I. Fund. Math., 166(1-2):1–82, 2000.
- [She03a] Saharon Shelah. More Jonsson algebras. Arch. Math. Logic, 42(1):1–44, 2003.
- $\begin{array}{ll} \mbox{[She03b]} & \mbox{Saharon Shelah. Not collapsing cardinals} \leq \kappa \mbox{ in } (<\kappa) \mbox{-support iterations}. Israel Journal of Mathematics, 136:29-115, 2003. math.LO/9707225. \end{array}$
- [She03c] Saharon Shelah. Successor of singulars: combinatorics and not collapsing cardinals $\leq \kappa$ in ($< \kappa$)-support iterations. Israel Journal of Mathematics, 134:127–155, 2003. math.LO/9808140.
- [She04] Saharon Shelah. Forcing axiom failure for any $\lambda > \aleph_1$. Arch. Math. Logic, 43(3):285–295, 2004.
- [She08] Saharon Shelah. Reflection implies the SCH. Fund. Math., 198(2):95–111, 2008.
- [She10] Saharon Shelah. Diamonds. Proc. Amer. Math. Soc., 138(6):2151–2161, 2010.
- [She13] S. Shelah. On incompactness for chromatic number of graphs. Acta Math. Hungar., 139(4):363–371, 2013.
- [She21] Saharon Shelah. Divide and conquer: dividing lines and universality. *Theoria*, 87(2):259–348, 2021.
- [SheXX] Saharon Shelah. Non-structure theory. Oxford University Press, XX. to appear.
- [Shi99] Masahiro Shioya. Splitting $P_{\kappa}\lambda$ into maximally many stationary sets. Israel J. Math., 114:347–357, 1999.
- [Shi08] Masahiro Shioya. Diamonds on $P_{\kappa}\lambda$. In Computational prospects of infinity. Part II. Presented talks, volume 15 of Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., pages 271–298. World Sci. Publ., Hackensack, NJ, 2008.
- [SS10] Daniel Soukup and Lajos Soukup. Club guessing for dummies, 2010.
- [SZ02] Saharon Shelah and Jindřich Zapletal. Duality and the PCF theory. Math. Res. Lett., 9(5-6):585–595, 2002.
- [Tod02] Stevo Todorčević. Localized reflection and fragments of PFA. In Set theory (Piscataway, NJ, 1999), volume 58 of DIMACS Ser. Discrete Math. Theoret. Comput. Sci., pages 135–148. Amer. Math. Soc., Providence, RI, 2002.
- [Tod07] Stevo Todorcevic. Walks on ordinals and their characteristics, volume 263 of Progress in Mathematics. Birkhäuser Verlag, Basel, 2007.
- [Tod18] Stevo Todorcevic. Collapsing ω_2 with semi-proper forcing. Arch. Math. Logic, 57(1-2):185–194, 2018.
- [Zap01] Jindřich Zapletal. Transfinite open-point games. Topology Appl., 111(3):289–297, 2001.
- [Zap04] Jindřich Zapletal. Descriptive set theory and definable forcing. Mem. Amer. Math. Soc., 167(793):viii+141, 2004.

[Zap08] Jindřich Zapletal. Forcing idealized, volume 174 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2008.

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT-GAN 5290002, ISRAEL.

Department of Mathematics, Bar-Ilan University, Ramat-Gan 5290002, Israel. $\mathit{URL:}\ \mathtt{http://www.assafrinot.com}$